

**RISK FACTORS ASSOCIATED WITH OVERWEIGHT AND  
OBESITY AMONG REPRODUCTIVE AGED FEMALE  
RESIDING IN ITAHARI SUB-METROPOLITAN CITY**

by

**Samir khulal**

**Department of Nutrition and Dietetics**

**Central Campus of Technology, Dharan**

**Institute of Science and Technology**

**Tribhuvan University, Nepal**

**2022**

**Risk factors Associated with Overweight and Obesity Among  
Reproductive Aged Females Residing in Itahari Sub-metropolitan City**

*A dissertation submitted to Department of Nutrition and Dietetics, Central Campus of  
Technology, Tribhuvan University in partial fulfillment of the requirements for the  
Bachelors degree of Nutrition and Dietetics.*

by

**Samir Khulal**

**T.U. Registration No. 5-2-8-86-2016**

**Department of Nutrition and Dietetics**

**Central Campus of Technology, Dharan**

**Institute of Science and Technology**

**Tribhuvan University, Nepal**

**November, 2022**

**Tribhuvan University**  
**Institute of Science and Technology**  
**Department of Nutrition and Dietetics**  
**Central Campus of Technology, Dharan**

**Approval Letter**

This *dissertation* entitled *Risk Factors Associated with Overweight and Obesity Among Reproductive Aged Females Residing in Itahari Sub-metropolitan City*, presented by **Samir khulal** has been accepted as the partial fulfillment of the requirement for the degree of **Bachelors of Science in Nutrition and Dietetics**.

**Dissertation Committee**

**1. Head of the Department**

**(Asst. Prof. Kabindra Bhattari, HOD)**

**2. External Examiner**

**(Mr. Dinesh Subedi, Food Reaserch Officer)**

**3. Supervisor**

**(Mrs. Pallavi Vyas Jaisani, Teaching Asst.)**

**4. Internal Examiner**

**(Mr. Suman Pokhrel, Teaching Assistant)**

**December, 2022**

## **Acknowledgements**

In the first place, I would like to express my sincere gratitude to my supervisor Mrs. Pallavi Vyas Jaisini, Teaching Assistant, for her continuous support throughout my graduate study. Her patience, motivation, enthusiasm and immense knowledge have guided me in research and writing of the thesis study. She has also consistently been motivating and helpful as I have progressed academically.

Secondly, I owe my sincere gratitude to Asst. Prof. Dambar Bahadur Khadka, former Department Head and Asst. Prof. Kabindra Bhattarai, Department Head of Nutrition and Dietetics, Central Campus of Technology for their valuable advice on my research work and providing with all necessary facilities for dissertation. I would also like to thank Campus Chief of Central Campus of Technology, Dr. Dil Kumar Limbu. This dissertation work was a great learning experience for me and has played a pivotal role for directing my academic career into right course.

Also, I would like to take this opportunity to thank my seniors Ms. Smriti Chauhan and Samikshya Pradhan, who have guided me throughout my dissertation period and to my friends Samir Karki, Sagar Khatiwoda, Binod Khanal, Rishiraj Adhikari, Vatsala Rai who gave me their time and thoughts during my research work. I would also like to thank my juniors for their time in my work and all the participants of this study. I owe a debt of gratitude to the Itahari Sub-Metropolitan City and all the participating members who agreed to participate in the interview and gave me the time and information I required.

At last, I owe my deepest gratitude to my respected parents who shed their sweat for making me “me” what I am now.

Date of submission: December, 2022

---

(Samir Khulal)

## **Abstract**

The study was intended to assess risk factors associated with overweight and obesity among Reproductive Aged Females residing in Itahari Sub-metropolitan city. A cross-sectional qualitative study was performed on 202 reproductive-aged females of 15-49 years of age with a structured questionnaire. Weight, height, waist and hip circumference were measured to determine indicators of overweight and obesity. General obesity was determined using WHO BMI classification while, WC and WHR analyzed abdominal obesity as per IDF and WHO criteria respectively. For data analysis, Microsoft package 16 (Excel and Word) and SPSS Statistics version 20 was used. Chi-square test was used to establish association between variables under study.

The study revealed 29.2% of respondents were overweight and 9.4% were obese; 59.9% of them were abdominally obese by WC and 67.8% by WHR. Here, mean BMI was  $24.20 \pm 3.91$  kg/m<sup>2</sup> and mean WC was  $85.55 \pm 10.53$  cm with mean WHR of  $0.90 \pm 0.06$  cm. Factors like age, marital status, parity, sleeping hours, eating pattern/vegetarianism, alcohol, egg and fast foods were factors found significantly associated ( $p < 0.05$ ) with overweight and obesity. Similarly, Age, marital status, education, occupation, family size, parity, contraceptive and physical adequacy were associated ( $p < 0.05$ ) with abdominal obesity. However, Factors like consumption of alcoholic drinks, egg and processed foods were only significantly associated ( $p < 0.05$ ) with BMI. Factors like, occupation and physical adequacy were only significantly associated ( $p < 0.05$ ) associated with WC. The study depicted the prevalence of overweight and obesity among reproductive aged females residing in Itahari sub-metropolitan city. Thus, overweight and obesity should be viewed as serious problems/issue.

## Contents

---

<b>Approval Letter .....</b>	<b>iii</b>
<b>Acknowledgements .....</b>	<b>iv</b>
<b>Abstract .....</b>	<b>v</b>
<b>List of tables .....</b>	<b>x</b>
<b>List of figures .....</b>	<b>xi</b>
<b>List of Plates .....</b>	<b>xii</b>
<b>List of abbreviations .....</b>	<b>xiii</b>
<b>Introduction .....</b>	<b>1-5</b>
1.1 General introduction .....	1
1.2 Statement of problem .....	2
1.3 Conceptual framework .....	4
1.4 Objectives .....	5
1.4.1 General objectives .....	5
1.4.2 Specific objectives .....	5
1.5 Research questions .....	5
1.6 Significance .....	5
1.7 Limitations.....	5
<b>Literature review .....</b>	<b>6-29</b>
2.1 Overweight and obesity .....	6
2.2 Measurement of overweight and obesity .....	7
2.2.1 Body mass index.....	7
2.2.2 Fat percentage.....	9
2.2.3 Waist circumference .....	10
2.2.4 Waist to hip ratio .....	11
2.3 Types of obesity .....	12

2.3.1 Based on BMI.....	12
2.3.2 Based on age of Onset of obesity .....	12
2.3.3 Based on distribution of Fat storage.....	13
2.4 Theories on obesity.....	15
2.5 Risk factors associated with overweight and obesity .....	16
2.5.1 Genetics .....	16
2.5.2 Age .....	17
2.5.3 Marital status .....	17
2.5.4 Socioeconomic factors.....	18
2.5.5 Physical activity.....	18
2.5.6 Dietary intake .....	21
2.5.7 Behavioral factors.....	24
2.6 Comorbidities of overweight and obesity.....	26
2.7 Prevalence and trends of overweight and obesity .....	27
2.7.1 Global trends of overweight and obesity .....	27
2.7.2 Overweight and obesity in Nepal .....	28
<b>Materials and methods.....</b>	<b>30-36</b>
3.1 Study area .....	30
3.2 Study population.....	30
3.3 Selection criteria .....	30
3.3.1 Inclusion criteria .....	30
3.3.2 Exclusion criteria.....	30
3.4 Research Design .....	30
3.5 Sampling technique .....	30
3.6 Sample size .....	31
3.7 Research instruments.....	32
3.8 Study variables .....	32

3.8.1	Independent variables .....	32
3.8.2	Dependent variables .....	33
3.9	Pre-testing.....	33
3.10	Validity and reliability.....	33
3.11	Data collection techniques.....	34
3.12	Data analysis.....	35
3.13	Logistic and Ethical considerations.....	36
<b>Result and Discussion.....</b>		<b>37-57</b>
4.1	Demographic and Socio-economic characteristics.....	37
4.1.1	Age wise distribution.....	37
4.1.2	Religion and caste.....	37
4.1.3	Marital status .....	38
4.1.4	Socio-economic factors .....	39
4.1.5	Family type and family size.....	40
4.1.6	Family history of obesity.....	40
4.2	Behavioral characteristics.....	41
4.3	Physical activity pattern .....	43
4.4	Health related factors.....	44
4.5	Dietary intake .....	45
4.5.1	Dietary characteristics .....	45
4.5.2	Dietary intake in preceding day.....	45
4.5.3	Food consumption pattern .....	47
4.6	Prevalence of overweight and obesity .....	49
4.6.1	According to international BMI classification .....	49
4.6.2	According to Asian BMI cut off.....	50
4.6.3	According to waist circumference.....	51
4.6.4	According to waist-to-hip ratio measurement .....	52



4.7 Factors associated with overweight and obesity.....	53
4.7.1 Factors associated with BMI (WHO cut off).....	53
4.7.2 Factors associated with waist circumference.....	56
4.7.3 Factors associated with waist-to-hip ratio .....	58
<b>Conclusions and recommendations.....</b>	<b>59</b>
5.1 Conclusions .....	59
5.2 Recommendations .....	59
<b>Summary .....</b>	<b>61</b>
<b>References.....</b>	<b>62-77</b>
<b>Appendices.....</b>	<b>78-87</b>

## List of tables

<b>Table no.</b>	<b>Title</b>	<b>Page no.</b>
2.1	WHO BMI classification	8
2.2	Classification of Asian BMI cut-offs	8
2.3	Percent body fat charts for male and female	10
2.4	WHO waist circumference cut-offs in male and female	11
2.5	MET values computation	20
4.1	Distribution of age in the surveyed population	37
4.2	Distribution of religion and caste of surveyed population	38
4.3	Distribution of marital status of surveyed population	39
4.4	Distribution of socio-economic factors of surveyed population	40
4.5	Distribution of family type and family size	41
4.6	Distribution of family history of obesity	41
4.7	Distribution of behavioral characteristics	42
4.8	Distribution physical activity pattern of surveyed population	44
4.9	Distribution of health-related factors of surveyed population	44
4.10	Distribution of dietary characteristics	45
4.11	Distribution of dietary intake in preceding day	46
4.12	Distribution of food consumption pattern of surveyed population	48
4.13	Factors associated with overweight and obesity based on BMI of WHO cut-off	54
4.14	Factors associated with abdominal obesity of surveyed population	57
4.15	Factors associated with abdominal obesity based on WHR	59

## List of figures

<b>Figure no.</b>	<b>Title</b>	<b>Page no.</b>
1.1	Conceptual framework for overweight and obesity	4
2.1	Trends of overweight and obesity in Nepal	29
4.1	Prevalence of overweight and obesity among reproductive aged female residing in Itahari sub-Metropolitan city	50
4.2	Prevalence of overweight and obesity by Asian BMI cut-offs	51
4.3	Prevalence of obesity with respect to waist circumference	52
4.4	Prevalence of obesity with respect to waist -to-hip ratio	53

## List of Plates

<b>Plate No.</b>	<b>Title</b>	<b>Page no.</b>
P1	Map of Itahari Sub-metropolitan city	91
P2	Measurement of height	92
P3	Measurement of weight	92

## List of abbreviations

<b>Abbreviation</b>	<b>Full form</b>
ADIPOQ	Adiponectin
BIA	Bioelectrical Impedance
BMI	Body Mass Index
CBS	Central Bureau of Statistics
CDC	Centers for Disease Control and Prevention
CHD	Coronary Heart Disease
CKD	Chronic Kidney Disease
CT	Computed Tomography
CVD	Cardiovascular Disease
DEXA	Dual-energy X-ray Absorptiometry
DFTQC	Department of Food Technology and Quality Control
FAO	Food and Agriculture Organization
FFQ	Food Frequency Questionnaire
GON	Government of Nepal
HC	Hip Circumference
HDL	High Density Lipoprotein
ICMR	Indian Council of Medical Research
IDF	International Diabetes Federation
IPAQ	International Physical Activity Questionnaire
LEPR	Leptin Receptor
MET	Metabolic Equivalent
MetS	Metabolic Syndrome
MOHP	Ministry of Health and Population
MRI	Magnetic Resonance Imaging
NCD	Non-communicable Disease
NDHS	Nepal Demographic and Health Survey
PA	Physical Activity

RDA	Recommended Dietary Allowances
SPSS	Statistical Package for Social Sciences
ST	Skinfold Thickness
T2DM	Type 2 Diabetes Mellitus
UNICEF	United Nations Children's Fund
WC	Waist Circumference
WOF	World Obesity Federation
WHO	World Health Organization
WHtR	Waist-to-Height Ratio
WHR	Waist-to-Hip Ratio

---

# Part I

## Introduction

### 1.1 General introduction

Overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health (WHO, 2022). Body mass index (BMI) is a simple index of weight-for-height that is commonly used to classify overweight and obesity in adults. It is defined as person's weight in kilograms divided by square of its height in meters ( $\text{kg}/\text{m}^2$ ). For adults, overweight is BMI greater than or equal to  $25 \text{ kg}/\text{m}^2$  and obesity is BMI greater than or equal to  $30 \text{ kg}/\text{m}^2$  (WHO, 2022). The fundamental cause of obesity and overweight is an energy imbalance between calories consumed and calories expended. An increased intake of energy dense food and increasingly sedentary nature of life style and work, and increasing urbanization have led to global epidemic of overweight and obesity (WHO, 2021).

Body mass index (BMI) is a simple index of weight-for-height that is commonly used to classify overweight and obesity in adults. It is defined as a person's weight in kilograms divided by the square of his height in meters ( $\text{kg}/\text{m}^2$ ) (Shahi *et al.*, 2013b). In 2015, 41% of CVD-related death, 9.5% of diabetes-related death and 10% of death from CKD and neoplasms were attributed to obesity (Afshin *et al.*, 2017). Compared with adults of normal weight, adults with BMI  $>40 \text{ kg}/\text{m}^2$  have an approximately 64% higher risk of T2DM, a 54% higher risk of high blood pressure, a 9% higher risk for high cholesterol, a 17% higher risk for asthma, a 34% higher risk for arthritis, and a 32% higher risk for generally fair or poor health (Must *et al.*, 2012).

Recently, evidence has been growing to suggest that an additional indicator of central obesity, waist circumference (WC), may be more closely associated with metabolic risks compared to BMI because it reflects regional body fat distribution and upper body adiposity (Sahakyan *et al.*, 2015).

In 2016, more than 1.9 billion (39%) adults aged 18 years and older were overweight and of these over 650 million (13%) adults were obese. The overweight population consists of 39% men and 40% women whereas the obese population consists of 11% men and 15% women. The worldwide prevalence of obesity nearly tripled between 1975 and 2016 (WHO,

2022). In developing countries with under-nutrition, obesity is a complex condition, with serious social and psychological dimensions, affecting virtually all ages and socio-economic groups (WHO, 2004).

In case of Nepal, it was found that 24.3% of adult population were overweight or obese which consists of 23.4% of male and 25.1% of female population. Also, percentage of adults who were obese was 4.3% including 3.2% of male and 5.3% of female. The mean waist to hip ratio of female was found to be 0.90. 63.6% adults (56.3% male and 70.2% female) have high prevalence of WHR (Dhimal *et al.*, 2020).

Itahari is a sub-metropolitan city in the Sunsari District of Province No. 1 of Nepal. Itahari city is business hub of eastern Nepal. It is the second most populous city in Eastern Nepal after Biratnagar. Situated at a distance of 25 kilometres north of the provincial capital of Biratnagar, 16 kilometres south of Dharan and 92 kilometres west of Kakarbhitta, Itahari serves as a junction point of the east-west Mahendra highway and the north–south Koshi highway. Itahari has an estimated city population of 1,98,098 living in 40,207 households as per 2021 Nepal census.

## **1.2 Statement of problem**

Nepal's increasing trend towards urbanization and leads health challenges, whose consequences are seen as overweight and obesity. Changing dietary habits can shift a society's disease pattern from infectious, communicable diseases' dominance towards a status of double-disease burden with increasing prevalence of obesity and non-communicable diseases (Vaidya *et al.*, 2010). Overweight and obesity is a significant public health concern affecting more than half a billion people worldwide (Gakidou, 2014).

Nepal is experiencing nutrition transition in recent decades because of which there is increase consumption of high fat and high sugar foods. Rapidly growing economies of the country has resulted in the globalization of food markets, fast food chains and the increasing availability of street vendors who offer products at very competitive value due to economical acquisition of inputs such as raw and processed foods which increases the consumption of energy dense food (Bhurosy and Jeewon, 2014).


The obesity burden would certainly cost heavy in low- and middle-income countries. Many low and middle income countries have been neglecting overweight and obesity as



major health treats, with policies in place to tackle undernutrition, but lack policies to halt the given burden of diseases due to rise of overweight and obesity (WHO, 2013). An enabling environment is of paramount importance to increase awareness about the risk factors for overweight and obesity in general population to decrease the prevalence of overweight associated NCDs in the upcoming generations of Nepal (Piryani *et al.*, 2016b). It is the responsibility of policy makers and all the other concerned authorities to address this issue.

Looking at the urbanization rate in Itahari sub-metropolitan city and the observed high prevalence of risk factors in adults it becomes necessary to assess the nutritional status of adults to find out over nutritional status. Thus, assessment of overweight and obesity in male and female is must needed in order to know the prevalence of overweight and obesity along with its risk factors. Such an assessment will help policy maker and developer to address the fast-growing problem appropriately and in a timely manner to reduce the chronic health impact of overweight and obesity among male and female as well as associated consequences of NCDs in the upcoming generations of Nepal.

### 1.3 Conceptual framework



**Figure 1.1** Conceptual framework for overweight and obesity

(Jiménez, 2013; Sartorius *et al.*, 2015)

## **1.4 Objectives**

### **1.4.1 General objectives**

The general objective of the study was to identify the prevalence of overweight and obesity among reproductive aged females of 15-49 years residing in Itahari sub-metropolitan city.

### **1.4.2 Specific objectives**

- i. To assess overweight and obesity among reproductive aged females of 15-49 years residing in Itahari sub-metropolitan city.
- ii. To assess physical activity level, behavioral factors and dietary factors with the help of questionnaire.
- iii. To identify risk factors directly and indirectly associated with overweight and obesity.

## **1.5 Research questions**

- i. What is the prevalence of overweight and obesity among reproductive aged females living in Itahari sub-metropolitan city?
- ii. What are the risk factors associated with the prevalent overweight and obesity among them?

## **1.6 Significance**

- i. This study will be helpful in highlighting the distribution of overweight and obesity and its associated contributing factors.
- ii. The result of this study will be helpful for the adults to improve their dietary habits and physical activity level.
- iii. As health problems associated with obesity and overweight are increasing more often now a day; these findings will be helpful in informing the health sector and the public health planners in mobilization and allocation of resources for the prevention and control of NCDs.

## **1.7 Limitations**

- i. Obesity was not assessed by the body fat percentage due to limited resources.

## PART II

### Literature review

#### 2.1 Overweight and obesity

Obesity is a medical condition described as excess body weight in the form of fat. Overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health (WHO, 2022). Body mass index (BMI) is a simple index of weight-for-height that is commonly used to classify overweight and obesity in adults. It is defined as person's weight in kilograms divided by square of its height in meters ( $\text{kg}/\text{m}^2$ ). For adults, overweight is BMI greater than or equal to  $25 \text{ kg}/\text{m}^2$  and obesity is BMI greater than or equal to  $30 \text{ kg}/\text{m}^2$  (WHO, 2022). BMI is a measure of generalized obesity whereas central obesity can be measured on the basis of waist circumference and waist to hip ratio (WHR). An adult who has a BMI of 25-29.9 is considered overweight, and an adult who has a BMI over 30 is considered obese. A BMI of (18.5-24.9)  $\text{kg}/\text{m}^2$  is considered normal weight. In general, overweight and obesity indicate a weight greater than what is considered healthy. Obesity is a chronic condition defined by an excess amount of body fat. According to WHO waist to hip ratio above 0.90 for male and 0.85 for female is considered as central obesity whereas WC above 90 cm for male and 80 cm for female is considered as being centrally or abdominally obese (IDF., 2005).

When energy intake equals energy expenditure, the body is in positive energy balance and body energy is stable. Over the many years, when body remains in positive energy balance, that is, the intake of calories is more than the expenditure of energy, this leads to obesity. The generalized accumulation of excess adipose tissue in the body with body weight more than 20% of the desirable body weight is called obesity whereas overweight is body weight 10-20% more than the mean standard weight for age, height and sex (Srilakshmi, 2019). Obesity is a complex issue related to lifestyle, environment, and genes. Environmental and genetic factors have a complex interaction with psychologic, cultural, and physiologic influences (Mahan and Raymond, 2016).

Obesity is characterized by high cholesterol and fatty acid levels, imbalance in metabolic energy, insulin desensitization, lethargy, gallstones, high blood pressure, shortness of breath, emotional and social problems, and excessive adipose mass accumulation with hyperplasia and hypertrophy (Kushner and Ryan, 2014).

When energy intake equals energy expenditure, the body is in positive energy balance and body energy is stable. Over the many years, when body remains in positive energy balance, that is, the intake of calories is more than the expenditure of energy, this leads to obesity. The generalized accumulation of excess adipose tissue in the body with body weight more than 20% of the desirable body weight is called obesity whereas overweight is body weight 10-20% more than the mean standard weight for age, height and sex (Srilakshmi, 2019). The positive energy balance is a broad aspect for increase in adiposity; there are various internal factors to be blamed. Neurochemicals, body fat stores, protein mass, hormones, and post-ingestion factors, all play their roles in regulating intake and weight maintenance. Short term regulation governs hunger, appetite and satiety related to consumption of food from meal to meal and long term regulation of body weight is controlled by the availability of adipose stores and hormone responses involving insulin and leptin (Mahan and Raymond, 2017).

## **2.2 Measurement of overweight and obesity**

Body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR) and waist-height ratio (WHR) are different measures used to estimate the body fat content; however, comparison to a gold standard test is practically difficult (Baïoumi, 2019). The most accurate measures of body weight and composition are underwater weighing, dual-energy x-ray absorptiometry (DEXA) scanning, computed tomography (CT), and magnetic resonance imaging (MRI) but are impractical for use in everyday clinical encounters as well as in research settings (Purnell, 2018). The practicable, inexpensive and routine measurement includes anthropometric measurement of body weight, height, waist and hip circumference required to construct indicators of adiposity namely, body mass index (BMI), Waist circumference and Waist-to-hip ratio (Kushner, 2012).

### **2.2.1 Body mass index**

Body Mass Index (BMI), also known as Quetlet Index, is a measure combining the weight and height of an individual. It can be calculated by dividing the weight(in kg) by the square of the height (WHO, 2022).

The BMI cut-offs are given by WHO and are categorized as follows:

**Table 2.1: WHO BMI Classification**

<b>Classification</b>	<b>BMI (kg/m<sup>2</sup>)</b>	<b>Risk of comorbidities</b>
Underweight	<18.5	Low
Normal	18.5-24.9	Average
Overweight/ Pre-obese	25-29.9	Increased
Obese I	30-34.9	Moderate
Obese II	35-39.9	Severe
Obese III	≥40	Very severe

However due to high body fat content in Asians, they develop negative health consequences at a lower BMI than western populations. South-Asians have an increased body fat percentage (BF %), both total and in the abdominal region, lesser lean mass, skeletal muscle and bone mineral content along with a higher risk for CVD. The significant variability in body composition between ethnic groups may not be truly reflected by measuring only BMI or other markers as each has its own limitations. Therefore, WHO recommended lower cut-off points of BMI for high risk populations including South Asians (Amin *et al.*, 2015).

**Table 2.2: Classification of Asian BMI cut-offs**

<b>Classification</b>	<b>BMI(kg/m<sup>2</sup>)</b>	<b>Risk of comorbidities</b>
Underweight	<18.5	Low risk
Normal	18.5-23	Acceptable risk
Overweight	23-27.5	Increased risk
Obese	≥27.5	High risk

BMI provides the most useful population level measure of overweight and obesity as it is the same for both sexes and for all ages of adults. However, it should be considered a rough guide because it may not correspond to the same degree of fatness in different individuals (WHO, 2022). It measures excess body weight for a particular height and has been shown to

correlate with body fat although it is not a direct measure of body fat. It does not measure overweight or obesity risk and mortality risk with the same accuracy in all target populations due to variations in body fat composition and distribution (Bhurosy and Jeewon, 2014).

### **2.2.2 Fat percentage**

Body fat is composed of cells that contain triglyceride stores. Body fat can be divided into two main compartments: subcutaneous and intra-abdominal. Fat may also accumulate at ectopic sites such as muscle, including the heart, pancreas, and liver (Depres and Lemieux, 2006; Ronti *et al.*, 2006). Body fat percentage is total body fat expressed as a percentage of total body weight. The generally accepted definition of obesity, based on total body fat, uses cut-points as >25% in men, and >30% in women (Okorodudu *et al.*, 2010). Percent body fat provides a more accurate body assessment for being overweight or obese in compared with BMI (Trang *et al.*, 2019). Excessive ectopic fat distribution in sites such as the viscera leads to abnormal fat metabolism and aggravates the development of diabetes and obesity (Hwang *et al.*, 2015).

The determination of the accuracy of any anthropometric measure to determine body fat is gauged by the method of fat identification to which it is compared as all these anthropometric measures are indirect measures of fat distribution. The direct methods are those which have the ability to identify fat droplets all over the body with particular relevance to visceral fat and ectopic fat, for example, in skeletal muscle, liver, pancreas, heart (Baïoumi, 2019).

Computed Tomography (CT) or dual-energy X-ray absorptiometry (DEXA), continue to be the gold standard for evaluating the distribution of body fat (Rollins *et al.*, 2017). Nevertheless, the high cost and low availability have made it difficult to use in large population studies. Recently, more research has examined the potential role of body composition measurements in health monitoring (Mazzocoli, 2016; Ramirez-Velez *et al.*, 2017). Bioelectrical impedance analysis (BIA) is the method that is most frequently used to assess body composition and calculate body fat composition and calculate body fat percent in clinical practice, given its accuracy, simplicity, low cost, and excellent correlation with DEXA, CT, or magnetic resonance imaging (MRI) (Xiaoguang *et al.*, 2011). In addition to them, body fat percentage estimated by skin fold thickness (ST) is widely practiced indirect measure in health researches (Sangachin *et al.*, 2018). According to age the adjusted body fat percentage of women can be categorized as follows:

**Table 2.3:** Percent body fat charts for woman

<b>Age in years</b>	<b>Low Fat</b>	<b>Normal</b>	<b>Overweight</b>	<b>Obese</b>
<b>20-39</b>	<21%	21-32.9%	33-38.9%	≥39%
<b>41-60</b>	<23%	23-33.9%	34-39.9%	≥40%
<b>60-79</b>	<24%	24-35.9%	36-41.9%	≥42%

(Gallagher *et al.*, 2000)

### **2.2.3 Waist circumference**

Waist circumference is an indicator of health risk associated with excess fat around the waist. It is an anthropometric measure at waist line that provides a simple and practical measure for assessing central adiposity. It has been shown that mesenteric adipose tissue inflammation is more related to metabolic consequences of obesity (Kranendonk *et al.*, 2015). Waist circumference has been increasingly used as a convenient measure of adipose tissue in epidemiological studies as well as in weight loss intervention trials (Filgueiras *et al.*, 2019). It has been suggested that waist circumference is more practical and superior to BMI in case of predicting obesity-related mortality (Jacqui, 2017). This is largely based on the rationale that increased visceral adipose tissue is associated with a range of metabolic abnormalities, including decreased glucose intolerance, reduced insulin sensitivity and adverse lipid profiles, which are risk factors for type 2 DM, cancers and CVD (Seidell, 2010).

Waist circumference is measured at the midpoint between the lowest palpable rib and top of iliac crest, using a non-stretchable measuring tape that provides a constant 100g tension; at the end of several consecutive breaths (WHO, 2011). The WHO stated cut offs for waist circumference and risk of metabolic complications are as follows:



**Table 2.4:** WHO waist circumference cut-offs in male and female

<b>Classification</b>	<b>Cut-offs</b>	<b>Metabolic risk</b>
<b>Centrally overweight</b>	>80 cm	Increased
<b>Centrally obese</b>	>88 cm	Substantially increased

(WHO, 2011)

However, the recommended cut off values of waist circumference for central obesity vary among different ethnic groups. Asians tend to have more body fat per BMI than Caucasians, which indicates greater potential for them to develop hypertension, diabetes, and dyslipidemia at lower BMIs (Ma *et al.*, 2013). Therefore, the recommended cut off values for defining abdominal obesity among South Asians are  $\geq 90$ cm for males and  $\geq 80$  cm for females (IDF, 2006).

#### **2.2.4 Waist to hip ratio**

The waist-to-hip ratio (WHR) is obtained by dividing the waist circumference by the hip circumference (HC) using the same units of measurements for both. The HC is measured using a non-elastic tape held horizontally without constricting it at the point that yields the maximum diameter over the buttocks (Baïoumi, 2019). For the WC and the HC, each measurement should be done twice and if both measurements differ by >1cm, each measurement should be repeated (WHO, 2005). The role of WHR as a surrogate measure of both visceral and subcutaneous fat was established following convincing evidence from longitudinal studies that showed significant associations between abdominal obesity and adverse cardiovascular outcomes (Umuerrri, 2019).

Abdominal obesity is defined as waist-to-hip ratio above 0.90 for males and above 0.85 for females. These values indicate abdominal obesity and reflect substantially increased cardio-metabolic risks (WHO, 2011). The ratios are defined as one of the decisive benchmarks for metabolic syndrome and is consistent with findings of research predicting all cause and cardiovascular disease mortality (Mahan and Raymond, 2017).

Measuring hip circumference may be more difficult than measuring waist circumference alone; this could limit the potential use of waist hip ratio as an alternative to either waist circumference alone or BMI (WHO, 2011). WHR can also be harder to interpret than waist

circumference and have little implications with those who have a BMI of 35 or higher (Watson and Bubnis, 2017).

## **2.3 Types of obesity**

### **2.3.1 Based on BMI**

Based on increasing BMI, obesity can be categorized into following three different grades (Srilakshmi, 2019):

i. Grade I obesity

People with this type of obesity have body mass index more than 25 kg/m<sup>2</sup> but less than 29.9 kg/m<sup>2</sup>. Overweight does not affect their health; they may lead normal health and life expectancy is above normal. They may reduce on their own (Srilakshmi, 2019).

ii. Grade II obesity

The body mass index is between (30-39.9) kg/m<sup>2</sup>. Though they appear to be in good health, they have reduced tolerance to exercise with shortness of breath on exertion and they are unduly fatigued. This is due to the burden of increased weight they always carry and have reduced capacity of the circulatory and respiratory systems that are handicapped by masses of internal fat and fatty infiltration of muscle. For metabolic and mechanical reasons these patients are at increased risk of diabetes, atherosclerosis, hypertension, fatty liver, gall bladder diseases, osteoarthritis, hernias and varicose veins. Mortality rate also increases (Srilakshmi, 2019).

iii. Grade III obesity

The body mass index is above 40 and these patients are in pathetic conditions. Their day-to-day activities are restricted due to their enormous mass and more susceptible to diseases mentioned in Grade II. They are susceptible to atherosclerosis, prone to accidents and have serious psychological disturbances (Srilakshmi, 2019).

### **2.3.2 Based on age of Onset of obesity**

Patients with early onset of obesity had a higher total body fat mass, and higher body fat percentage, and a 1.84 times higher risk of BMI above 40 kg/m<sup>2</sup> than patients with adult onset of obesity ( $\geq 20$  years) (Wrzosek and Wisniewska, 2018). On the basis of age of onset, obesity can be categorized into two ways (Srilakshmi, 2019):

i. Juvenile-onset obesity

Juvenile obesity occurs due to hyperplasia and most rapidly in the first few years of life. It appears most frequently in the first year of life, at 5-6 years of age and during adolescence (Srilakshmi, 2019). Too many calories fed in infancy and early childhood leads to an overproduction of fat cells (hyperplasia) followed by hypertrophy (enlargement of fat cells) (Sheth and Shah, 2006). Overweight and obese children are likely to stay obese into adulthood and more likely to develop non-communicable disease like diabetes and cardiovascular diseases at a younger age (Sahoo *et al.*, 2015). Childhood obesity can profoundly affect children's physical health, social, and emotional well-being, and self-esteem. It is also associated with poor academic performance and a lower quality of life experienced by the child. Many co-morbid conditions like metabolic, cardiovascular, orthopedic, neurological, hepatic, pulmonary, sleep apnea and renal disorders are also seen in association with childhood obesity (Sahoo *et al.*, 2015).

ii. Adult -onset obesity

It occurs from hypertrophy of fat cells alone; the size of the individual cell is greatly enlarged. A distended adipose cell leads to further physiological, biochemical, anatomic aberrations in individual's organs and organ systems. In addition to this fact, increased number of fat cells may also occur in adult life which is to be expected when BMI is above 40 kg/m<sup>2</sup> (Bray and Bouchard, 2019). A pharmacology intervention coupled with other management strategies will work better in treating such cases than juvenile obesity (Srilakshmi, 2019). High energy density diet, increased portion size, low physical activity and adoption of a sedentary lifestyle as well as eating disorders are considered as important risk factors for the development of obesity (James, 2008). These behavioral and environmental factors lead to alterations in adipose tissue structure (hypertrophy and hyperplasia of adipocytes, inflammation) and secretion (e.g. adipokines) leading to development of metabolic syndrome (Tilg and Moschen, 2006).

### **2.3.3 Based on distribution of Fat storage**

Fat mass is distributed differently in men and women. The pattern of body's adipose tissue deposition, independent of total body fat, alters health risk in human body despite of their age and gender. There are two types of regional fat distribution as follows (McArdle *et al.*, 2010):

i. Android-type obesity

In central or android-type obesity, there is increased health risk from fat deposition, particularly internal visceral deposits, resulting from active lipolysis from the adipose tissue with catecholamine stimulation. Fat stored in this region shows greater metabolic responsiveness than fat deposited below waist line; as reflected by altered metabolic profile such as insulin resistance, hyperinsulinemia, non-insulin dependent diabetes mellitus, hyperlipidemia and hypertension (McArdle *et al.*, 2010).

Android obesity is commonly related to male body and is also called as apple-type body however, this can be related to menopause in case of females. The underlying mechanism might be due to the lipolytically sensitive abdominal depots providing excess free fatty acids to muscle tissues that has a decreased capacity for their oxidation. This excessive exposure of fatty acids impairs the insulin function of tissues and increase blood insulin levels leading to reduced insulin sensitivity and ultimately making a person insulin resistant. The hyperinsulinemia might be the starting point for most of other metabolic derangements (Srilakshmi, 2019). Also, increase in central fat more readily support processes that cause heart disease (McArdle *et al.*, 2010).

ii. Gynoid-type obesity

Gynoid-type obesity is related to weight gain around hips and flank area; thigh and bottom. It is typically related to obesity in women and is also known as pear-type or peripheral fat distribution (Srilakshmi, 2019). Individuals with a gynoid fat distribution are at a greater risk of mechanical problems like osteoarthritis (Sheth and Shah, 2006). The lower body fat deposition is usually by hyperplasia. Reducing the number of fat cells in lower hyperplastic depot is difficult than android obesity. This explains the weight loss difficulties faced by many women with gynoid obesity (Srilakshmi, 2019).

Gynoid fat deposition in women during child bearing years is utilized to support the demands of pregnancy and lactation. Women with the gynoid-type obesity do not develop the impairments of glucose metabolism as compared with an android deposition (L.K. Mahan and Raymond, 2017).

## 2.4 Theories on obesity

Obesity arises in a person when the size or the number of fat cells increases in the body. A normal person has got about 30-35 billion fat cells (Sheth and Shah, 2006). When a person tries to lose fat, then it is possible to reduce the size of fat cells rather than their numbers. Hence with initial increase in number of fat cells, weight loss becomes difficult (Hopkin, 2008). Different theories have been postulated to understand the body weight regulation mechanism and subsequent obesity. These include:

### i. Fat cell theory

The number of fat cells are determined early in life which once have formed, have a tendency to form full fat. The fat cells increase early in life (juvenile-onset hyperplasia) which indicates that adult-onset obesity is caused by an increase in the size of fat cells (hypertrophy). When a person starts losing weight, the cells decrease in size but the number generally stays the same. Thus, people having large number of fat cells have more difficulty in maintaining body weight than those with fewer fat cells (Srilakshmi, 2019). However, the obesity related health problems occur due to the enlarged fat cells which are metabolically active rather than the number of fat cells or the person's weight. The number of fat cells increase as a result of positive energy balance but can decrease only with sustained weight loss for a prolong period of time (Spalding *et al.*, 2008).

### ii. Set point theory

According to the set point theory, body weight remains remarkably stable from the internal regulatory mechanisms that are genetically determined. A deliberate efforts to starve or overfeed are followed by a rapid return to the original body weight called a 'set point' (L.K. Mahan and Raymond, 2017). Thus, once a body weight reaches this point, a whole set of signals is produced that influences the person's food intake to maintain in set point (Srilakshmi, 2019).

The set point mechanism is not regulated on a meal-to-meal basis. During the recovery from illness, food intake is increased in a catch-up fashion until lost weight is regained. Similarly, when the weight of the person exceeds the set point, then the hypothalamus recognizes the need to lose weight (Sheth and Shah, 2006).

### iii. Enzyme and hormone theories

When there is hyperinsulinemia, then the lipogenesis takes place which leads to conversion of glucose into triglycerides (fat). The resulting high serum triglycerides are stored in fat cells i.e., in the adipose tissue, and make the fat cells distended. This defect of chemical imbalance has been described as syndrome “X”. It makes you fat and becomes difficult to lose weight (Sheth and Shah, 2006).

When the fat cells absorb more of triglycerides, they emit a biochemical protein into bloodstream called leptin. When there is too much of leptin, it signals to restrict feeding behavior; the fat storage is inhibited and stimulates lipolysis with corresponding energy expenditure. Hence, leptin is regarded as a bodyweight regulatory hormone (Srilakshmi, 2019). In obesity, though leptin levels remain elevated, their receptors in the hypothalamus are desensitized. So, the hypothalamus continues the triglyceride storage in adipose cells with reflex of hunger beyond need and ultimately results in weight gain. Also, an increase in lipoprotein lipase enzyme is known to deposit fat into fat cells and has probable role in raising the appetite (Sheth and Shah, 2006).

## **2.5 Risk factors associated with overweight and obesity**

Obesity is a complex multifactorial chronic disease developing from interactive influences of numerous factors including - social, behavioral, psychological, metabolic, cellular and genetic. The various influencing factors on energy intake and expenditure that are considered to be important in weight gain and the development of obesity are as follows (L.K. Mahan and Raymond, 2017):

### **2.5.1 Genetics**

Obesity tends to run in families. It is likely that some individuals are genetically more susceptible to the effects of an obesogenic environment (Gandy *et al.*, 2006). Many hormonal and neural factors involved in weight regulation are determined by heredity and genetics. Small defects in their expression or interaction could contribute significantly to weight gain. Genetic inheritance influences 50-70 percentage a person’s chance of becoming fat more than any other factor (L.K. Mahan and Raymond, 2017). Within a family, the chance of being obese is 80 percent if both parents are obese and 50 percent if one parent is

obese. A mutation in the human gene coding for the B3 receptor in adipose tissue, involved in lipolysis and thermogenesis markedly increase the risk of obesity (Srilakshmi, 2019).

Genes interact with the diet and activity patterns that lead to obesity and the metabolic pathways that influence satiety and energy balance. Researchers have identified an obesity gene called 'Ob' that is expressed primarily in the adipose tissue and codes for the protein leptin (Whitney and Rolfes, 2011). Although numerous genes are involved, several have received much attention- the 'Ob' gene, the adiponectin (ADIPOQ) gene, the "fat mass and obesity associated" gene or FTO gene, and the beta3-adrenoreceptor gene. The Ob gene produces leptin. Mutation in the Ob gene, leptin receptor (LEPR), or ADIPOQ genes can result in obesity or metabolic syndrome (MetS) (Mahan and Raymond, 2017).

### **2.5.2 Age**

The aging process brings about many changes in body composition, often without concomitant changes in body weight and body mass index. In general as individual's age increases, percentage body fat increases whereas lean mass and bone mineral density decrease (Choi *et al.*, 2012). Advancing age is associated with an increase in abdominal white adipose tissue and fat deposition in skeletal muscle, due to hormonal changes and a less active lifestyle which may increase the risk of obesity with significantly affecting insulin sensitivity (Jura and Kozak, 2016). The body fat distribution also changes during aging, such as increased fat accumulation from the subcutaneous area to visceral organs (Öztürk *et al.*, 2018). Hales *et al.* (2018) found that age standardized prevalence of severe obesity in adults increased from 5.7% (95%-CI) in (2007-2008) to 7.7% (95%-CI) in (2015-2016). The prevalence increased in men, women and adults aged 20 to 39 years and 40-59 years.

### **2.5.3 Marital status**

Marital status affects body weight. In particular, living without partner, either being divorced or never married is associated with lower body weight. Cohabitors and married respondents tend to weigh more. There are three perspectives linking body weight to marital status (Teachman, 2016). The first perspective, the resource model, emphasizes different resources, social and economic, available to individuals possessing different marital status. Married individuals are more likely to have a confident with whom to eat and may therefore eat more

regularly and of larger portion sizes, as well as decreased physical activity, leading to weight gain (Dinour *et al.*, 2012).

The second model, the attractiveness model, links body weight to differences in emphasis people place on their physical attractiveness. Married men and women are less likely to be conscious or concerned about their body weight because they are not actively seeking a mate. As a consequence, married individuals are more likely to experience greater increase in body weight when compared to non-married individuals (Wilson, 2012). The third model is the crisis model and it focuses on stresses associated with marital status, particularly marital dissolution. Stress linked to marital disruption have been related to psychological, physiological and social consequences that can lead to weight loss (Teachman, 2016).

#### **2.5.4 Socioeconomic factors**

Socioeconomic status is an important factor associated with obesity. It can be determined using variables such as education, income, and occupation, with education considered to be the most stable variable over time (Anekwe *et al.*, 2020). The prevalence of obesity varies with socioeconomic status. In developed countries, poverty is associated with increase prevalence, whereas in developing countries it is the relative affluence that carries the greater risk (Yeung and Laquatra, 2003). The growing urbanization and globalization of food production and marketing, changing socioeconomic status of population has a profound effect on energy balance with introduction of low-cost, energy-dense foods in the domestic food market (Bhurosy and Jeewon, 2014).

The specific factors such as social identity, social status, social trends and influences of the built, industrial, and social environments are closely associated with the prevalence or incidence of obesity or that impacts efforts to prevent and treat this disease (Lee *et al.*, 2000). Socioeconomic factors contribute to obesity on an individual and community level, and any viable approach to sustainably addressing the obesity epidemic must take these factors into account (Anekwe *et al.*, 2020).

#### **2.5.5 Physical activity**

Lack of physical activity, through its impact on energy balance, has been identified as an important modifiable risk factor for obesity (Bryan and Walsh, 2004). There has been increasing evidence over the past decades of the importance of physical exercise in



maintaining cardiovascular health and preventing diseases (Blair and Moriss, 2009). When sedentary individuals undertake exercise, the activity provides a massive stimulus with widespread physiological implications. The precise metabolic regulation brought about by exercise is expressed at many levels of regulatory processes, be it by stimulating the effect of key enzymes, by increasing cell sensitivity to numerous hormones, by facilitating substrate transport through membranes, by influencing cell receptors in a tissue-specific manner, and much more. With the generalized sedentariness observed in modern societies, the human body needs to compensate for the lack of exercise stimulation to maintain energy and macronutrient balance. Fat gain and the metabolic syndrome are unfortunately the price to pay to maintain this balance (Chaput *et al.*, 2011).

Most of the research clearly report the prevalence of overweight and obesity being higher among the individuals with low levels of physical activity as compared to those with high levels of physical activities (Dabrowska *et al.*, 2015). However, the physical activity interventions had a positive effect on adiposity measures (Baker *et al.*, 2016). It is recommended that adults aged 18-64 years should do at least 150 minutes of moderate intensity aerobic physical activity throughout the week, or do at least 75 minutes of vigorous intensity aerobic physical activity throughout the week, or an equivalent combination of moderate and vigorous intensity physical activity (WHO, 2010). Also, the intensity of physical activity is measured in metabolic equivalents or METs. One MET is defined as the calories burned while an individual sit quietly for one minute. For the average adult, this is about one calorie per every 2.2 pounds of body weight per hour (IPAQ, 2005).

One of the methods for the assessment of physical activity level in community level is ‘International Physical Activity Questionnaire (IPAQ)’. From the questionnaire, total MET minutes/week and physical activity level can be determined as following (Ashok *et al.*, 2016; IPAQ, 2005):

**Table 2.5:** MET values computation

<b>MET values</b>	<b>Formula for computation</b>
Walking MET minutes/week	$3.3 \times \text{walking minutes} \times \text{walking days}$
Moderate MET minutes/week	$4 \times \text{moderate intensity activity minutes} \times \text{moderate days}$
Vigorous MET minutes/week	$8 \times \text{vigorous intensity activity minutes} \times \text{vigorous days}$
Total MET minutes/week	Walking + Moderate + Vigorous MET minutes/week scores

(Ashok *et al.*, 2016; IPAQ, 2005)

After the calculation of total MET score of each participant, the physical activity level can be categorized according to IPAQ scoring protocol as follows:

**i. Low:**

This is the lowest level of physical activity. Those individuals who do not meet criteria for moderate and high level or with MET scores less than 600 MET minutes/week are considered of having low level of physical activity.

**ii. Moderate:**

The pattern of activity having following criteria can be classified as moderate:

- 3 or more days of vigorous intensity activity of at least 20 minutes per day. OR
  - 5 or more days of moderate intensity activity and/or walking of at least 30 minutes per day.
- OR
- 5 or more days of any combination of walking, moderate intensity or vigorous intensity activities achieving a minimum total physical activity of at least 600 MET minutes/week.

**iii. High:**

Individuals meeting following criteria can be classified of having high physical activity level:

- Vigorous intensity activity on at least 3 days achieving a minimum total physical activity of at least 1500 MET minutes/week. OR
- 7 or more days of any combination of walking, moderate intensity or vigorous intensity activities achieving a minimum total physical activity of at least 3000 MET minutes/week.

Moderate intensity physical activity is defined as activities that are strenuous enough to burn three to six times as much energy per minute when sitting quietly, or 3 to 6 METs. Vigorous intensity activities burn more than 6 METs.

### **2.5.6 Dietary intake**

Diet plays an important role in obesity. Excess calorie intake along with overconsumption of specific nutrients and foods such as saturated fat, added sugars, high sodium food, and refined grain contribute to obesity (Jensen *et al.*, 2014). Carbohydrates, fat and protein all contribute to the body's energy needs. To meet the need for these macronutrients without increasing risk of chronic disease requires that people eat a balance diet. The recommendations for population nutrient intake goals include 55-75% total carbohydrate intake as adequate, total fat consumption of 15-30% and protein ingestion of 10-15% (WHO/FAO, 2010).

#### **i. Energy dense foods**

Energy density is the amount of energy (in kilocalories) per given weight of food (in grams). Energy dense foods, often high in refined grains, added sugars, and added fats (Bechthold, 2014), are palatable, inexpensive, and convenient (Mendoza *et al.*, 2007). A single group of researchers have identified dietary fats, sugars and corn sweeteners, soft drinks, fast foods, snacks, and larger portion sizes as potential culprits in the obesity epidemic (Drewnowski, 2007). Dietary fat has higher energy density (9 kcal/g) than other macronutrients; either carbohydrate or protein (4 kcal/g), and excess dietary fat is readily stored in adipose tissue depots with a very high efficiency. Disruptions in fat balance has become a clear perspective of weight changes following challenges to body weight regulation (WHO, 2003).

Lower food costs may be associated with more energy-dense diets, and total energy intake may actually increase. Lower food costs may be associated with more energy-dense diets, and total energy intake may actually increase (Drewnowski, 2009). The etiology of obesity increasingly reflects excessive calorie intake matched with higher levels of sedentary activity that occur in the face of a worldwide urban migration. In this scenario, traditional diets are often replaced with low cost energy dense foodstuffs produced by the industrialized food (Gluckman *et al.*, 2015; Hill and Peters, 1998). Nepalese diet is shifting away from agricultural staple foods to modern processed foods with higher total energy, total fat, and

sugar resulting in increased prevalence of overweight/obesity and diet related non-communicable diseases (Subedi *et al.*, 2017).

#### ii. Protein intake

The word protein is derived from the Greek word “proteios”, meaning prime or primary. This term is very appropriate in nutrition, because protein is the most fundamental component of tissues in animals and human. A protein contains various amount of different amino acid linked via peptide bonds (Wu, 2016). An adequate level of protein consumption is essential for every individual, and the World Health Organization (WHO) has established an international dietary protein recommendation of 0.83 g/kg/day (WHO, 2015).

Protein has long been regarded for its ability to assist in weight management, increase high-density lipoprotein (HDL) cholesterol, increase thermogenesis and satiety, and enhance bone mineralization (Ahnen *et al.*, 2019). In the 7 year follow up study, Bujnowski *et al.* (2011) reported that animal protein intake was positively related and vegetable protein intake inversely associated with overweight/obesity in apparently healthy middle-aged American men. The results of this study indicate that protein sources may be important with respect to weight, independent of energy, carbohydrate, alcohol, and fat intake (Bujnowski *et al.*, 2011). A cross-sectional study conducted in multiethnic population concluded that high dietary protein intake protects against the effects of risk variants in the FTO gene on BMI and waist circumference (Merritt *et al.*, 2018).

#### iii. Fruits and vegetables

Fruits and vegetables have historically held a place in dietary guidance because of their concentrations of vitamins, especially vitamins A and C; minerals, especially electrolytes; and more recently phytochemicals, especially antioxidants. They are also good source of dietary fibers (Slavin and Lloyd, 2012). It is suggested to consume more than 400 grams (five servings) of fruits and vegetables per day to improve overall health and reduce the risk of certain NCDs excluding tubers like potato, cassava, etc (WHO, 2020). In Nepal, low consumption of fruits and vegetables was found which can be directly attributed to prevailing non-communicable diseases (Shahi *et al.*, 2013a).

Consequently, replacing some added sugars and saturated fat with more fruits and vegetables may benefit health. Some research indicates that fruit and vegetable consumption

reduces the risk of major diseases and possibly delays the onset of age-related disorders (Vincente *et al.*, 2014). Fruits and vegetables are high in water and fiber, incorporating them in the diet can reduce energy density, promote satiety and decreased energy intake (Rolls *et al.*, 2004). Research shows that some bioactive components in fruits and vegetables, such as carotenoids, could contribute to the prevention and treatment of obesity. Some of these carotenoids are responsible for vitamin A production, a hormone-like vitamin with pleiotropic effects in mammals. Among these effects, vitamin A is a potent regulator of adipose tissue development, and is therefore important for obesity (Coronel *et al.*, 2019).

#### iv. Milk and Milk products

Milk products play an important role in daily human nutrition, as milk contains a balanced composition of amino acids, vitamins, and minerals (Smolnikova *et al.*, 2021). It is a source of anti-oxidative peptides, Vitamins, organic acids, calcium and pro-biotic bacteria with other biologically active peptides that are liberated from milk proteins by different hydrolytic methods. Besides milk, many dairy products including butter, yoghurt, cream, cheese, and kefir, have been manufactured and consumed worldwide (Patil *et al.*, 2021). Milk and dairy products constitute the best dietary source of calcium due to the bioavailability of the calcium they contain (Rovira, 2015).

Greater consumption of total dairy products may be of importance in the prevention of weight gain in middle aged and elderly women who are initially normal weight. Epidemiologic data suggest that people with greater dairy product (high calcium) intake have a lower prevalence of overweight and obesity (Rautiainen *et al.*, 2016). The possible anti-obesity mechanisms of calcium includes regulation of adipogenesis, fat metabolism, adipocyte (precursor) proliferation and apoptosis, thermogenesis, fat absorption and excretion, and gut microbiota (Zhang *et al.*, 2019).

#### V. Salt intake

The recommended salt intake for adults is less than 5 gram of salt per day which is 2 grams of sodium per day (WHO, 2012). High sodium intake is significantly associated with an increased risk of obesity and metabolic syndrome in the general population (Oh *et al.*, 2015). Recently, several lines of evidence have also shown that high salt intake is associated with an increased risk of obesity. One reason for this association is that high salt intake stimulates

thirst and increases fluid intake and thereby increasing sugar-sweetened beverage consumption (Ma *et al.*, 2015).

Sodium intake also stimulates appetite, and subsequently increases energy intake and extracellular volume (Oh *et al.*, 2017). In addition, the association between salt and obesity may also be partially caused by excessive consumption of processed food that is high in both calorie and salt (Ma *et al.*, 2015). A diet with a high salt content may have a negative influence on development in body composition by expanding body fat and reducing fat free mass (Larsen *et al.*, 2013).

### **2.5.7 Behavioral factors**

The behavioral influences on obesity include specific behaviors that may contribute to weight gain through overeating or reduced physical activity and the mechanisms by which the environment can affect health (Affenito *et al.*, 2012)

#### **i. Watching T.V. while eating**

Television watching appears to encourage snacking during viewing and also influences food choices both during viewing and at other times. In controlled interventions, decreased television watching reduced weight gain in children, and effect that was mediated more by improvements in dietary habits than by a change in physical activity (Mozaffarian *et al.*, 2011). The association between watching television with overweight and obesity was evidenced by a different explanation such as hours spent in front of the television, decrease in the physical activity, a television based food advertising influencing what and how people eat, and watching television leads to mindless eating or lack of attention paid to consume due to external cues in the environment (Ahmed *et al.*, 2020).

The more television people watch, the more likely they are to gain weight or become overweight or obese (Thorp *et al.*, 2011). Different studies revealed that watching television is positively associated with an overall increase in food intake, particularly pizza, fast food, and high-calorie snacks and is inversely associated with intakes of vegetables and fruits (Ahmed *et al.*, 2020). A research among adolescents in Nepal, reports that watching TV for longer time is one of the major risk factors for developing overweight (Piryan *et al.*, 2016b).

## ii. Skipping breakfast

“Breakfast is the most important meal of the day” is a consensus in public. However, the definition of breakfast varies across studies. Here are two standard definitions as follows: Firstly, Eating the first meal of the day before or at the beginning of the activity within 2 hours of waking up, usually no later than 10 am, the calorie content is 20% to 35% of the total daily energy requirement. Secondly, the consumption of food or beverage (excluding water) between 5 and 9 am (Ma *et al.*, 2020).

Breakfast skipping is associated with changes in appetite and decreased satiety, which may lead to subsequent overeating and impaired insulin sensitivity (Önnerfält *et al.*, 2018). Chowdhury *et al.* (2016) conducted a randomized controlled trial and demonstrated that in obese adults, daily breakfast leads to more significant physical activity during the morning, whereas morning fasting results in partial dietary compensation (i.e. higher energy intake) later in the day. Skipping breakfast have become more prevalent among school-age children, adolescents and working adults (Ma *et al.*, 2020). Skipping breakfast was significantly correlated with waist circumference and BMI (Watanabe *et al.*, 2014).

## iii. Sleep

Sleep plays a crucial role in human’s endocrine, metabolic and neurologic functions (Liu *et al.*, 2019). Sleep is an important modulator of neuroendocrine function and glucose metabolism and sleep loss has been shown to result in metabolic and endocrine alterations, including decreased glucose tolerance, decreased insulin sensitivity, increased evening concentrations of cortisol, increased levels of ghrelin, decreased levels of leptin, and increased hunger and appetite (Beccuti and Pannain, 2011). Experimental sleep restriction was associated with increased salt retention and inflammatory markers as well. Many epidemiological studies have shown that decreased sleep duration and quality (<7 hours per night) is significantly associated with increased obesity incidence (Cooper *et al.*, 2018). Thus, quantity and quality of sleep can also prevent a risk factor for overweight and obesity through dysfunctional eating behaviors, decreased physical activity, and metabolic changes (Bonanno *et al.*, 2019).

iv. Eating outside once in a day

Eating outside may lead to overconsumption and increase risk of obesity in part because of larger portion sizes, high energy dense foods, and increased variety and preferred taste of the foods (Anderson *et al.*, 2011). Food choices when eating out are usually high in energy content, which contributes to excessive energy intake (Bezerra *et al.*, 2012). Studies have found a positive association between the intake of food away from home and person's body mass index (BMI) or weight gain (Seguin *et al.*, 2016). Even one-meal per week consumption outside home, in fast food and sit-down restaurant, was associated with increase in BMI (Bhutani *et al.*, 2016). Fast foods typically have an extremely high energy density, which leads to excessive energy intake that contributes to weight gain and obesity (Prentice and Jebb, 2003).

v. Alcohol intake

Alcohol drinking is a component of lifestyle in many regions and, recently, has been raised in some developed countries. It contains high energy (7.1 kcal/g) and also, by disrupting energy balance, leads to accumulation of fat mass and development of overweight or obesity (Golzarand *et al.*, 2021). Increased energy intake through alcohol consumption can promote an energy imbalance, where intake exceeds output, and ultimately contribute to weight gain if not compensated for (Traversy and Chaput, 2015).

Regular heavy episodic drinking in young adulthood is associated with higher risk of gaining excess weight and transitioning to overweight/obesity (Fazzino *et al.*, 2017). Light-to-moderate alcohol intake, especially wine, may be more likely to protect against weight gain, whereas consumption of spirits has been positively associated with weight gain (Carmen *et al.*, 2011). Alcohol has numerous complex effects on, for instance, food intake regulation, psychosocial well-being, sleeping, and depression symptoms, all of which may influence body weight (Traversy and Chaput, 2015). Beer intake > 500 mL/day has been associated with abdominal obesity ("beer belly") (Bendsen *et al.*, 2013).

## **2.6 Comorbidities of overweight and obesity**

Understanding the pathophysiology of obesity related disease will allow providers to better manage the suite of diseases that affect an individual. Keeping obesity as the central focus and basis of all treatment strategies for obesity related diseases is essential for the management of current disease as well as the prevention of future diseases (Andolfi and



Fisichella, 2018). Health consequences of obese people fall into two broad categories: those attributable to the effect of increased fat mass (such as osteoarthritis, obstructive sleep apnea, social stigmatization) and those due to the increased number of fat cells ( diabetes, cancer, cardiovascular disease, non-alcoholic fatty liver disease) (Gupta *et al.*, 2011).

Obesity may be associated with hypertension, dyslipidemia, diabetes, or insulin resistance, and elevated levels of fibrinogen and C-reactive protein, all of which increase the risk of CVD mortality and morbidity, particularly with central deposition of adipose tissues (Akhil and Ahmad, 2011). The relative risk of mortality from cancer, attributable to obesity, was approximately 14.2% in men and 19.8% in women. The most highly associated with obesity include breast cancer in postmenopausal women, colon cancer (especially among men), endometrial, esophageal adenocarcinoma, gall bladder and renal cancers (Stone *et al.*, 2018). The long-term risk of type 2 diabetes increases significantly with increasing weight. More than 6 fold increase in diabetes risk for class III obese (BMI $\geq$ 40) individuals was observed, compared to normal weight individuals (Leung *et al.*, 2017). In obesity, there is insulin resistance especially in muscle and there is hyperinsulinemia because of impaired insulin uptake by receptors in target tissue (Srilakshmi, 2019).

Obesity is characterized by a state of chronic and low-grade inflammation and obese patients are affected at various levels of their innate and adaptive immune responses. Obese patients have chronically lowered concentration of adiponectin (an anti-inflammatory adipokines) and higher levels of leptin (a pro-inflammatory adipokines) (Ouchi *et al.*, 2011). Obesity is a well-recognized cause of respiratory function compromise which might make these group of patients at risk of a more severe clinical course if they contract COVID-19. We have also concluded that obese subjects are potentially more vulnerable to become infected with COVID-19, and are more contagious when they do so with a prolonged period of viral shedding (Albashir, 2020). A rise in 10 kg body weight increases the risk of coronary artery disease by 12% and at the same time, systolic blood pressure rises by 3 mmHg and diastolic by 2.3 mmHg as a consequence (Csige *et al.*, 2018).

## **2.7 Prevalence and trends of overweight and obesity**

### **2.7.1 Global trends of overweight and obesity**

The prevalence of overweight and obesity has increased dramatically during the past four decades posing a major public health challenge both in developing and developed countries.

Worldwide obesity has nearly tripled since 1975 (WHO, 2022). Between 1975 and 2016, the prevalence of obesity increased in every country in the world. In 1995, there were an estimated 200 million obese adults worldwide which increased to over 300 million in 2000 (Agha and Agha, 2017). The obesity pandemic has continued to rise since then; it had reached 600 million in 2014 and over 650 million in 2016 (WHO, 2022). If these trends continue, by 2025, global obesity prevalence will reach 18% in men and surpass 21% in women (NCD-RisC, 2016). In 2016, more than 1.9 billion (39%) adults aged 18 years and older were overweight and of these over 650 million (13%) adults were obese. The overweight population consists of 39% men and 40% women whereas the obese population consists of 11% men and 15% women (WHO, 2022). Globally 8% of deaths in 2017 were the result of obesity which represents an increase from 4.5% in 1990 (Ritchie and Roser, 2020).


The prevalence of overweight has increased in almost all Asia-Pacific countries and territories since 2010. The increase was observed to be faster in lower –middle and low income countries and territories – 38% for children and adolescents (from 10.8% to 14.9%) and 15% for adults (from 26.9% to 30.8%) (OECD and WHO, 2020). Two out of every five adults are either overweight or obese in the Asia-Pacific regions (Helble and Francisco, 2017). The obesity burden is greater for South Asia, due to differences in fat patterning and body composition and the cardio-metabolic effects of body mass index (BMI) at levels far below standard BMI overweight cut-offs of 25 (Popkin *et al.*, 2012). Deaths attributed to obesity for South-Asian population increased from 2.03% in 2000 to 5.59% in 2017 (Ritchie and Roser, 2020).

### **2.7.2 Overweight and obesity in Nepal**

According to the STEPS Survey Nepal 2008, 7.2% adults were overweight or obese and 1.7% obese. The mean BMI was 20.6 kg/m<sup>2</sup> (MOHP *et al.*, 2009). The trend of overweight and obesity increased to 21.6% overweight or obese and 4% obese in 2013 with the mean BMI of 22.4 kg/m<sup>2</sup> (Aryal *et al.*, 2014). A survey in 2016 estimated that nearly 19.5% Nepalese adults were overweight or obese which includes 22% of female. The percentage of obese population was 3.5% (5% women and 2% men) (MOHP and ICF, 2017). According to the STEPS Survey Nepal 2019, the percentage of overweight or obese population was

24.3% (23.4% male and 25.1% female) and that of obese population was 4.3% (3.2% male and 5.3% female). The mean BMI was found to be 22.7 kg/m<sup>2</sup> (Dhimal *et al.*, 2020).

The prevalence of overweight and obesity was 22.9% in Province 1 and 3.8% of population were obese. Province 3 had the highest prevalence of overweight and obesity when compared with other provinces of Nepal (Dhimal *et al.*, 2020). A STEPS survey conducted in Nepal found that 7.2% and 1.7% of adults were overweight and obese respectively in 2007 which increased to 17.7% and 4% in 2013 respectively (MOPH, 2013). A study related Prevalence of Hypertension, Obesity, Diabetes, and Metabolic Syndrome in Nepal, the result found out that 28% were overweight, and 22% were obese (Sharma *et al.*, 2011a). Likewise another study done in Lalitpur sub metropolitan city described that almost 12.2% adolescents were overweight (Piryani *et al.*, 2016a). Overweight/obesity prevalence of 33.4% was found in a study conducted among civil servants in Nepal (Simkhada *et al.*, 2011). The study done in Ramdhuni municipality described that almost 26.9% and 14.3% adults residing in Ramdhuni municipality were overweight and obese respectively (Chaudhary, 2020).



(Aryal *et al.*, 2014; Dhimal *et al.*, 2020; MOHP and ICF, 2017; MOHP *et al.*, 2009)

**Figure 2.1:** Trends of overweight and obesity in Nepal

## **PART III**

### **Materials and methods**

#### **3.1 Study area**

The study was conducted at Itahari sub-metropolitan city of Sunsari district, Koshi zone, which is situated in Province no. 1 of Nepal. The total area of Itahari sub-metropolitan city was 93.79 sq km<sup>2</sup>. It has total population of 1,57,457 population of which 79965 are males and 77492 are females living in 50,743 households (CBS, 2021; Municipality, 2079).

#### **3.2 Study population**

The population under study were reproductive age females of 15-49 years of age residing in Itahari sub-metropolitan city.

#### **3.3 Selection criteria**

##### **3.3.1 Inclusion criteria**

15-49 years female residing in Itahari sub-metropolitan city were included in the study.

##### **3.3.2 Exclusion criteria**

- i. Respondents who were not available at household during the time of survey.
- ii. Those seriously ill, mentally unfit, pregnant and lactating their children.
- iii. Those who denied to give informed consent.

#### **3.4 Research Design**

A community based cross-sectional study was conducted among reproductive aged females from different wards of Itahari who were available at the time of field work. The field work consisted of survey with the help of structured questionnaire followed by anthropometric measurements.

#### **3.5 Sampling technique**

The study was conducted on 7 wards of Itahari sub-metropolitan city. Out of 20 wards in Itahari a total of 7 wards were selected for sample selection by using lottery method. Number of households to be surveyed from each ward was calculated on the basis of probability proportionate sampling technique to improve precision of sampling strategy. Random

households were chosen for sample selection. Only one female from each household were chosen for study.

### 3.6 Sample size

The sample size was determined by using a single proportional formula where the combined prevalence rate of overweight and obesity was taken to be 22% in the survey area (MOH, 2017), 95% confidence interval (CI), 6% margin of error (d) and 10% non-response rate is added to the total calculated sample size.

We have, the sample size for infinite population as:

$$\text{Sample size (N)} = Z^2 \times p(1-p)/d^2$$

where,

z = confidence level at 95% (standard value of z is 1.96)

p = baseline prevalence of overweight or obesity in Province 1

d = margin of error

$$\text{Now, } N = 1.96^2 \times 0.22 \times (1-0.22) / (0.06)^2 = 183.116 = 184$$

According to the population Census of 2079, the total female population of reproductive aged of Itahari was 47068. Thus, we apply finite population sample formula to obtain new sample size to conduct survey in Itahari.

Therefore,

$$\text{New SS} = N / [1 + (N-1) / \text{POP}]$$

Where,

New SS = New sample size for finite population

N = Sample size in infinite population

POP = Total number of female populations of reproductive aged

New sample size obtained as

$$= N / [1 + (N-1) / \text{POP}]$$

$$= 184 / [1 + ((184-1) / 47068)]$$

$$= 183.28 \approx 184$$

Thus, calculated sample size is adjusted for non-response. Considering non-response rate as 10%, the adjusted sample size is calculated to be 202.

### **3.7 Research instruments**

- i. Weighing machine: Weighing machine manufactured by Microlife Pvt. Ltd, with the capacity of 180kg and having the least count of 0.1Kg (1piece) was used.
- ii. Stadiometer: Stadiometer was used to measure height with the capacity of 197 cm and having the least count of 0.1cm.
- iii. Measuring tape: A non-stretchable flexible measuring tape was used to measure waist and hip circumference.
- iv. Questionnaire: A well designed, structured and pretested set of questionnaire was used to collect information on socio-demographic and economic data, physical activity, dietary intake and behavioral characteristics.
- v. Measuring utensils: Standardized utensils were used for taking 24 hour dietary recall.

### **3.8 Study variables**

#### **3.8.1 Independent variables**

- i. Socio-economic and demographic variables

Age, ethnicity, religion, marital status, education, occupation, family size.

- ii. Physical activity

Physical activity was classified into different levels as low, moderate and high according to the calculated score of each individual from the IPAQ-short questionnaire. Similarly, the adequacy of physical activity for each individual was also determined as per the global recommendations on physical activity for health (WHO, 2010).

- iii. Dietary intake

To assess the dietary intake, 24-hour dietary recall was used. The information collected was transformed to nutrient intake, namely, macronutrients. The nutrient adequacy was determined based on recommended dietary allowance (RDA) for Nepalese adults (DFTQC, 2017; ICMR, 2010). While salt intake (WHO, 2012) and fruits and vegetables intake (WHO, 2020) was based on WHO recommendations.

- iv. Behavioral characteristics

Watching TV while eating foods, skipping breakfast, smoking, alcohol intake, and sleep

- v. Health related characteristics

Use of contraceptives, menstrual disorders and thyroid problems.

### **3.8.2 Dependent variables**

- i. Body Mass Index (BMI): BMI is defined as person's weight in kilograms divided by square of its height in meters ( $\text{kg}/\text{m}^2$ ). Participants with a BMI greater or equal to 25 and less than  $30\text{kg}/\text{m}^2$  is considered as overweight and obese if BMI is greater than or equal to  $30\text{kg}/\text{m}^2$  (WHO, 2022).
- ii. Waist circumference: Female participants with waist circumference above 80 cm were identified as abdominally obese (Alberti *et al.*, 2006)
- iii. Waist to Hip ratio (WHR): Female respondents with WHR greater than 0.85 were considered as abdominally obese (WHO, 2011).

### **3.9 Pre-testing**

The set of questionnaires was prepared and reviewed by the supervisor. It was then pre-tested among 12 females for the practicability of the tool. This helped to have better understanding of the questions, estimation of the time consumed by each set of questions, and have clear view about the standardization procedures for anthropometry. After the completion of pretesting, all the misleading and wrongly interpreted questions were omitted and revised to obtain the final set of questionnaires.

### **3.10 Validity and reliability**

Validity refers to the degree to which the data collection instruments will measure what they intent to measure. Validity of weighing balance was ascertained by comparing the data provided by the weighing balance with the standard weights. Similarly, the validity of the stadiometer was ascertained by comparing the measurement with the UNICEF stadiometer. The measuring tape was calibrated against the standard stadiometer. The instruments were checked and reset daily to validate the data. For 24-hour recall, different foods were standardized in measuring cups used for data collection.

Reliability refers to the degree to which the result of a measurement, calculation, or the specification can be depended on to be accurate. It refers to how consistently a method measures something. The questionnaire was pre-tested prior to data collection to ascertain

the reliability and was checked daily for completeness and consistency. The test re-test method was used and close supervision was done in the field.

### **3.11 Data collection techniques**

Data collection was done on the month of October 2021, in two steps which consists of initial interview with the participants to complete the semi-structured questionnaire, followed by anthropometric assessment. An informed consent was obtained prior to data collection. Participants were asked about their general information, socio-demographic and economic information followed by anthropometry, physical activity level and dietary details.

i. Socio-demographic information:

The socio-demographic information involved asking the participants about their age, ethnicity, marital status, education, family annual income.

ii. Anthropometric assessment:

Each anthropometric measurement was repeated thrice.

- a. Weight: A portable digital weighing scale was used to measure weight. The instrument was placed on a firm, flat surface. Participants were requested to remove their footwear and socks, wear light clothes and stand in the scale with one foot on each side of the scale, face forward, place arms idly at their side and wait until asked to step off. Weight was measured in kilograms (Dhimal *et al.*, 2020; WHO, 2005)
- b. Height: A portable standardized stadiometer was used to measure height. The participants were asked to remove the footwear and any hat or hair ties. They were requested to stand on the flat board facing the interviewer, heels against the back board with their feet together and knees straight. They were asked to look straight ahead and not tilt their head up, making sure that their eyes are at the same level as ears. Height was measured in centimeters (Dhimal *et al.*, 2020; WHO, 2005)
- c. Waist circumference and Hip circumference: Waist circumference was measured at the mid-point between the lower margin of the least palpable rib and the top of the iliac crest, while, hip circumference was taken at the maximum over the buttocks; using a non-stretchable tape that provided a constant 100 g tension and at a level parallel to the floor. The tape was snugged around the body, but not pulled so tight that it was constricting. Also, the subjects were made stand erect with arms at the sides, feet positioned close together, and weight distribution across the feet and



relaxed with measurement taken at the end of normal respiration. The measurement was read at the level of the tape to the nearest 0.1 cm (Dhimal *et al.*, 2020; WHO, 2011)

iii. Physical activity:

Information on physical activity of the participants included its type, intensity, duration and frequency in a week for the purpose of work, transportation and recreational activities. The short form of 'International Physical Activity Questionnaire (IPAQ)' was used for the data collection. IPAQ short form is an instrument designed primarily for population surveillance of physical activity among adults (IPAQ, 2005). It is a valid instrument of measuring physical activity (PA) which allows the international comparisons and which studies PA in its totality (at work, at home, and in the context of transports and hobbies) (Ashok *et al.*, 2016).

iv. Dietary information:

Dietary assessment was done using a food frequency questionnaire (FFQ) and dietary recall (24-hour) method. The FFQ was used to estimate the usual diet pattern and understand the diet quality and obesity outcome based on the frequency of consumption of predetermined food list for a week. While, the 24-hour food recall involved asking participants to recall their food intake in the previous 24 hours (the previous day). This method assessed the food quantity in household standardized measure of measuring cups and table spoon. The gram equivalents of those foods were calculated which was used to calculate amount of nutrients consumed.

### **3.12 Data analysis**

The questionnaire was checked at the end of the day for completeness. The collected data sets were first coded and entered manually in database using Microsoft Excel 2016. Here, the qualitative data were transcribed and coded by assigning labels to various categories. The data was then transferred to IBM SPSS Statistics software (version 20) for further analysis. Descriptive analysis was used to describe percentage and distribution of respondents by socio-demographic variables, physical activity, dietary patterns, medical characteristics and behavioral characteristics. Chi square test was used to establish the relationships with explanatory variables in the assigned data set.

### **3.13 Logistic and Ethical considerations**

This research study was conducted with the permission received from the Department of Nutrition and Dietetics, Central Campus of Technology along with the ethical approval obtained from Itahari sub-metropolitan city. The objectives of research were explained in simple language and an informed written and verbal consent was obtained from all the participants. Privacy and confidentiality of collected data was ensured.

## Part IV

### Result and Discussion

The cross-sectional study to assess the prevalence of overweight and obesity with indicators like BMI, WC and WHR was conducted in 202 reproductive aged women from 15-49 years of age living in Itahari sub-metropolitan city. The following headings provide an overview of the findings:

#### 1.1 Demographic and Socio-economic characteristics

##### 4.1.1 Age wise distribution

The age group with the biggest population proportion was 21-30 years old, accounting for 33.2% (67), while the age group 15-20 years old had the lowest population share of 19.8% (40). The age distribution of study population is shown in Table 4.1:

**Table 4.1:** Distribution of age in the surveyed population (n=202)

Age group	Frequency	Percentage (%)
<20	40	19.8
21-30	67	33.2
31-40	51	25.2
41-49	44	21.8

##### 4.1.2 Religion and caste

Hindus comprised 94.6% (191) of the study's participants, followed by Buddhist and Muslim with equal population share i.e., 2% (4) and Christian with 1.5% (3). On the other side, the Chhetri group had the biggest percentage of participants 40.6% (82), followed by Janajati 29.2% (59); Brahmin 21.8% (44); Dalits 5.9% (12); and others 2.5% (5). The distribution of the study population by caste and religion is shown in Table 4.2 below:

**Table 4.2:** Distribution of religion and caste of surveyed population (n=202)

<b>Variable</b>	<b>Frequency</b>	<b>Percentage (%)</b>
<b>Religion</b>		
Hindu	191	94.6
Buddhist	4	2.0
Muslim	4	2.0
Christian	3	1.5
<b>Caste/Ethnicity</b>		
Brahmin	44	21.8
Chhetri	82	40.06
Dalits	12	5.9
Janajati	59	29.2
Others	5	2.5

#### **4.1.3 Marital status and parity**

Majority of the females, 68.8% (141) were married while the percentage of those widow/divorced were only 0.5% (1). Also, the study found that most of the females knew about the importance of family planning as 53.5% (108) had their children from 1 to 2 in numbers. 29.2% (59) were either unmarried, just married or planned their family later in life. However, 9.4% (19) had their parity of three or more. Table 4.3 shows how the study population's marital status and parity were distributed:

**Table 4.3:** Distribution of marital status of surveyed population (n=202)

<b>Variable</b>	<b>Frequency</b>	<b>Percentage (%)</b>
<b>Marital status</b>		
Unmarried	59	29.2
Married	141	69.8
Widow	1	0.5
separated	1	0.5
<b>Parity</b>		
0	75	37.1
1-2	108	53.5
≥3	19	9.4

#### **4.1.4 Socio-economic factors**

An individual's dietary preferences and consumption habits are influenced by their education level and gained information. According to the report, the majority of them, 84.2% (175) were literate, with the highest percentage of females with high school education, 30.2% (61). The recent finding from NDHS reported 33% of reproductive aged female are illiterate in Nepal (MOH, 2017). The comparative inference thus suggested females being more literate at itahari with only 15.8% (32) illiterate. Likewise, the type of occupation determines the income and level of physical activity performed by an individual during the work. More than half of the females, 72.8% (147) of reproductive age in itahari were economically dependent as they were unemployed; while rest others were 6.9% (14) daily wage workers, 7.4% (15) on job service, 5% (10) farmers, and 7.9 (16) were self-employed with their small business or shops. In addition to that, the distribution showed 55% (111) females had their monthly income below the average line while 45% (91) had higher income. The average monthly income defined for Nepalese is Rs. 30,121 (Nepal Rastra Bank, 2016). Table 4.4 illustrates the distribution of socioeconomic characteristics among the population sample.

**Table 4.4:** Distribution of socio-economic factors of surveyed population (n=202)

<b>Variables</b>	<b>Frequency(n)</b>	<b>Percentage (%)</b>
<b>Monthly income</b>		
<30000	111	55
≥30000	91	45
<b>Occupation</b>		
Daily wage worker	14	6.9
Service	15	7.4
Farming	10	5.0
Business	16	7.9
Unemployed	147	72.8
<b>Education</b>		
Illiterate	32	15.8
Primary school	29	14.4
Secondary school	56	27.7
High school	61	30.2
Graduate	24	11.9

#### **4.1.5 Family type and family size**

According to this study, the majority of participants, 83.7% (169) lived in nuclear families, while 16.3% (33) had joint families. The participants were shown to favor nuclear families. The result from recent national census found that average family size to be 4.32 (CBS, 2021). The study being consistent to it, found that the majority of the participants 54% (109) had a family size of less than five members. On the other hand, 46% (93) of the participants had larger family size. The distribution of type and size of family is shown in the table below:

**Table 4.5:** Distribution of family type and family size (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Type of family</b>		
Nuclear	169	83.7
Joint	33	16.3
<b>Family size</b>		
<5	109	54.0
≥5	93	46.0

#### **4.1.6 Family history of obesity**

When respondents were questioned about whether their parents had ever been overweight or obese; most of them, 40.6% (82) informed that their parents did not have gained excess weight while, 19.8% (40) reported both of them gained weight later in life, 25.7% (52) told only mother had excess weight and least, 13.9% (28) reported only their father was heavier weighed. The study population followed the following distribution as shown in Table 4.6:

**Table 4.6** Distribution of family history of obesity (n=202)

<b>Family history</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Both parents	40	19.8
Father only	28	13.9
Mother only	52	25.7
None	82	40.6

#### **4.2 Behavioral characteristics**

According to the survey, the vast majority of participants, 82.7% (167), did not watch television or use any other displays while eating any meals. On the other hand, 5.4% (11) watched three to four times each week, 5.9% (12) watched TV every day, and 5.9% (12) did so while eating meals twice every week.

Around 40.6% (82), skipped breakfast every day, whereas 21.3% (43) only did so twice or three times each week. No matter how hard they try, some of them claimed, they always forget to have breakfast once a week, as opposed to the 28.7% (58) who never skipped it. Similar to this, the survey found that 67.3% (136) of participants had an acceptable amount of sleep (7-9 hours) each night and rested, whereas 23.3% (47) slept less than 7 hours and 9.4% (19) slept more than 9 hours each night.

In the study, it was also found that the trend of eating-out of home seemed increasing as 30.7% (62) informed that they daily ate one or more meals out of their homes, 13.9% (28) often ate out when going shopping or any other purposes, 23.3% (47) ate out rarely and 32.2% (65) had never eaten out of home. Distribution of behavioral factors is given following in Table 4.8. Moreover, understanding the behavioral pattern of study population, the study revealed that 16.3% (33) used to drink alcohol either occasionally or on regular basis while 83.7% (169) did not drink alcoholic beverages. Also, regarding the smoking behavior, 98% (198) did not smoke tobacco but 2% (4) claimed about doing so. The distribution of these are presented in Table 4.7:

**Table 4.7:** Distribution of behavioral characteristics (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Eating in front of TV</b>		
Daily	12	5.9
Twice a week	12	5.9
3-4 times a week	11	5.4
Never	167	82.7
<b>Skipping breakfast</b>		
Daily	82	40.6
Once a week	19	9.4
Twice-Thrice a week	43	21.3
Never	58	28.7
<b>Sleep</b>		
<7 hours	47	23.3
7-9 hours	136	67.3



>9 hours	19	9.4
<b>Meals away from home</b>		
Daily	62	30.7
Often	28	13.9
Rarely	47	23.3
Never	65	32.2
<b>Alcohol intake</b>		
Yes	33	16.3
No	169	83.7
<b>Smoking/Tobacco</b>		
Yes	4	2.0
No	198	98.0

---

### 4.3 Physical activity pattern

According to the IPAQ scoring technique, which divided the individuals into three levels of physical activity, the participants' level of physical activity was evaluated. Low, moderate, and high levels of physical activity were the three categories. 52.5% (106) of the population were found to be physically inactive, followed by 45% (91) and 2.5% (5) who were physically active at moderate and high levels, respectively. According to an alternate analysis performed in accordance with WHO guidelines to determine the appropriate amount of physical activity, 47.5% (96) of the participants engage in adequate physical activity of more than 150 minutes per week while 52.5% (106) performed inadequate physical activity. Table 4.8 below displays how physical activity is distributed.

**Table 4.8:** Distribution of physical activity pattern of surveyed population (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Physical activity level</b>		
Low	106	52.5
Moderate	91	45.0
High	5	2.5
<b>Adequacy of physical activity</b>		
Inadequate	106	52.5
Adequate	96	47.5

#### **4.4 Health related factors**

It was found that 76.6% (155) females did not use any contraceptive measures in that cross-sectional period of study whereas others left, 23.3% (55) used any one of different contraceptives available at market for birth control. Likewise, the results showed that most female had good menstrual cycle as only 28.7% (58) suffered from menstrual irregularities while rest 71.3% (144) had monthly regular periods. The distribution of these factors is given below in Table 4.9:

**Table 4.9:** Distribution of health-related factors of surveyed population (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Contraceptive use</b>		
Yes	47	23.3
No	155	76.7
<b>Menstrual irregularities</b>		
Yes	58	28.7
No	144	71.3

## 4.5 Dietary intake

### 4.5.1 Dietary characteristics

According to the study, 97% (196) of the participants were not vegetarians. But among them, none of them adopted a vegan diet. However, 2.5% (5) a lacto-vegetarian diet, and 0.5% (1) a lacto-ovo vegetarian diet. Only 12.9% (26) of the participants had a salt intake that met WHO recommendations, according to their salt consumption patterns. 87.1% (176), consumed too much salt every day.

Majority of the participants, 50% (101) followed the consumption pattern of 3 meals per day followed by 42.1% (85) following the consumption pattern of 4 meals per day and 7.9% (16) had 5 meals per day. The distribution of dietary characteristics is shown in Table 4.10:

**Table 4.10:** Distribution of dietary characteristics (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Vegetarianisms</b>		
Vegan	0	0
Lacto-vegetarian	5	2.5
Lacto-ovo-vegetarian	1	.5
Non-vegetarian	196	97.0
<b>No. of meals</b>		
3	101	50.0
4	85	42.1
5	16	7.9

### 4.5.2 Dietary intake in preceding day

The result of the study showed that 32.2% (65) participants had consumed excess calories in their previous day. 43.6% (88) of them had inadequate calorie intake and only 24.3% (49) were found to have consumed adequate calorie. The mean calorie intake of the participants was found to be  $1863.31 \pm 197.88$  kilocalories which was lower than the minimum average

adequate requirement of 2220 kilocalories set by Government of Nepal (National Planning Commission & N. and Central Bureau of Statistics, 2013). The result was comparable with calorie intake of 1870 kcal/capita/day among urban households in Nepal (Kumar *et al.*, 2016).

91.6% (185) participants had adequate carbohydrate intake whereas 4.5% (9) of them had low carbohydrate intake and 4.0% (8) of them had high carbohydrate intake than the recommended amount. The mean carbohydrate intake was found to be  $296.81 \pm 48.40$  grams. This was lower than the mean consumption of  $404 \pm 100$  gm/day among rural population of southern Terai (Ohno *et al.*, 1997). The study revealed that 57.9% (119) participants had adequate intake of protein in their diet whereas 9.9% (20) had high intake of protein. Similarly, the mean protein intake was found to be  $58.96 \pm 13.93$  grams. The mean protein intake was greater than per capita protein intake of 51 gm/day in urban households of Nepal (Kumar *et al.*, 2016).

In case of fat consumption, it was found that 75.7% (153) of the participants had adequate intake of fat in their diet whereas 21.8% (44) had high fat intake. Also, 2.5% (5) of them had low fat intake. The mean fat intake was found to be  $48.90 \pm 13.70$  grams. This result was greater when compared with consumption of  $24.5 \pm 12$  gm/day in rural households of southern Terai region (Ohno *et al.*, 1997). The distribution of dietary intake of preceding day is shown in Table 4.11:

**Table 4.11:** Distribution of dietary intake in preceding day (n=202)

<b>Variable</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
<b>Calorie</b>		
Inadequate	88	43.6
Adequate	49	24.3
Excess	65	32.2
<b>Carbohydrates</b>		
Low	9	4.5
Adequate	185	91.6

High	8	4.0
<b>Protein</b>		
Low	65	32.2
Adequate	117	57.9
High	20	9.9
<b>Fat</b>		
Low	5	2.5
Adequate	153	75.7
High	44	21.8

---

#### 4.5.3 Food consumption pattern

Food consumption was classified as "regular" if it occurred at least once per day, "frequent" if it happened three to four times per week, "rare" if consumed once a week or less, and "never" if it didn't happen at all. It was discovered that all of the participants ate rice every day. This could be as a result of rice being our primary cereal grain. 29.7% (60) of the participants consumed wheat on regular basis and 34.2% (69) of them consumed wheat on frequent basis. Only 4% (8) of the participants consumed maize or millet or barley regularly and 5% (10) of them consumed on frequent basis, whereas the majority of them were discovered to just rarely or never consume them.

In the context of pulses and legumes, majority of the participants i.e., 47.5% (96) were found to have consumed polished daal on regular basis. Also, 25.7% (52) consumed polished daal respectively on frequent basis. Similarly, only 5.9% (12) consumed grams or beans or peas on regular basis while more than half of responded i.e., 55% (111) consumed these food items rarely.

In this study, 46% (93) and 88.1% (178) participants have claimed to have regular consumption of green leafy vegetables and other vegetables respectively. While 35.6% (72) and 4.9% (9) had consumed green leafy vegetables and other vegetables frequently respectively. In terms of fruits consumption, it was found that 27.2% (55) had regular consumption pattern and 49% (99) of them consumed frequently. However, 23.8% (48)

rarely consumed fruits. Comparing the results of fruits and vegetables consumption on daily basis, they were found higher than the daily consumption of fruits (14.5%) and vegetables (44.8%) among reproductive age women in terai belt of Nepal (Bhandari et al., 2016).

It was seen that 46.5% (94) consumed dairy products in regular basis in the forms like milk, milk tea, curd and butter milk mainly; 43.6% (88) frequently consumed them while 9.6% (14) consumed them rarely and 3% (6) never consumed. On comparison, the regular dairy products consumption in females of itahari was found to be higher than the consumption of females (34.1%) in overall terai region (Bhandari et al., 2016).

Among the animal sources, eggs were consumed more on regular basis 15.8% (32) whereas white meat and red meat were regularly consumed by 6.4% (13) and just 1% (2) of the respondents, respectively. The meat consumption pattern found chicken meat was more popular among the participants which comprises 42.6% (86) consuming it on frequent basis. Red meat was found to be less preferred by the participants. A large portion of study population 65.3% (132) had been found to consume red meat rarely and 24.8% (50) did not consume red meat at all. Similarly, intake of fast foods and processed foods, rich in calories, was assessed which concluded 27.2% (55) regularly consumed either fast food or processed foods, 46.5% (94) consumed on frequent basis while 24.8% (50) consumed them rarely. The distribution of food consumption pattern is shown in Table 4.12:

**Table 4.12:** Distribution of food consumption pattern of surveyed population (n=202)


<b>Variable</b>	<b>Regular</b>	<b>Frequent</b>	<b>Rarely</b>	<b>Never</b>
	<b>n(%)</b>	<b>n(%)</b>	<b>n(%)</b>	<b>n(%)</b>
<b>Rice</b>	202 (100)	0 (0)	0 (0)	0(0)
<b>Wheat</b>	60 (29.7)	69 (34.2)	73 (36.1)	-
<b>Maize/millet/barley</b>	8 (4.0)	10 (5.0)	149 (73.8)	35 (17.3)
<b>Polished daal</b>	96 (47.5)	52 (25.7)	35 (17.3)	19 (9.4)
<b>Grams/beans/peas</b>	12 (5.9)	79 (39.1)	111 (55)	-
<b>Green leafy vegetables</b>	93 (46.0)	72 (35.6)	37 (18.3)	-
<b>Other vegetables</b>	178 (88.1)	9 (4.5)	15 (7.4)	-

<b>Fruits</b>	55 (27.2)	99 (49.0)	48 (23.8)	-
<b>Dairy</b>	94 (46.5)	88 (43.6)	14 (6.9)	6 (3.0)
<b>Eggs</b>	32 (15.8)	69 (34.2)	78 (38.6)	23 (11.6)
<b>White meat</b>	13 (6.4)	86 (42.6)	84 (41.6)	19 (9.4)
<b>Red meat</b>	2 (1.0)	18 (8.9)	132 (65.3)	50 (24.8)
<b>Fast foods</b>	55 (27.2)	94 (46.5)	50 (24.8)	-
<b>Carbonated Beverages</b>	6 (2.9)	52 (25.7)	135 (66.8)	9 (4.6)

#### **4.6 Prevalence of overweight and obesity**

##### **4.6.1 According to international BMI classification**

The BMI result of the study was analyzed according to the international classification given by WHO which concluded that 58.4% (118) of them were normal, 29.2% (59) were overweight, 9.4% (19) were obese but only 3% (6) were underweight. Thus, the prevalence of overweight and obesity in Itahari Sub-metropolitan City was found much greater than the mean prevalence in Nepal of 24.3% (MOHP, 2020b); as well as share of overweight/obese (27.4%) in Province 1 (MOHP, 2017). Similarly, the result was higher than the recent STEPS survey report which claimed for overweight of 21.6% and obese of 3.8% in Province 1 ((MOHP *et al.*, 2020) . The prevalence of nutritional status among reproductive females in sub-metropolitan city of Itahari is depicted in Figure 4.1 below:



**Figure 4.1:** Prevalence of overweight and obesity among reproductive aged female residing in Itahari sub-Metropolitan city.


The mean BMI among study population was found to be  $24.20 \pm 3.91$  kg/m<sup>2</sup>; which was again higher than the recent average BMI of adult females (22.8 kg/m<sup>2</sup>) in Nepal (MOHP et al., 2020). Likewise, computing these figures against the survey done at western region of Nepal, the prevalence of overweight and obesity were 31.8% and 3.8% respectively (Koirala, 2019). The report clearly reflected overweight prevalence was very close but obesity proportion in Itahari sub-metropolitan city was about two times more. Similarly, it was found lower prevalence of 25% among women in Pakistan; though proportion of obese females (14%) was found to be higher than our achieved results (Janjua et al. et al., 2015)

#### 4.6.2 According to Asian BMI cut off

The result based on Asian BMI cut-off found that only 36.9% (80) were normal with 36. % (73) overweight, 21.3% (43) obese and 3% (6) underweight. The result was higher than the mean prevalence of overweight and obesity among women of 32.87%, according to Asian BMI cut-off, in Nepal (Rawal, 2018).

In another study, in terms of the Asian-specific BMI cutoffs, found the prevalence of overweight and obesity as 25.6% and 13.3% respectively among females all over Nepal, which again, was lower than our obtained results (Kibria, 2019). Figure 4.2 illustrates the distribution of overweight and obesity prevalence in Itahari sub-metropolitan city.






**Figure 4.2:** Prevalence of overweight and obesity by Asian BMI cut-offs

#### 4.6.3 According to waist circumference

The mean waist circumference was found to be  $85.55 \pm 10.53$  cm; higher than the population mean WC of all Nepalese adults i.e., 79.7 cm) (MOHP et al., 2020a). Likewise, mean WC in females of Chitwan was found to be  $85.96 \pm 7.112$  cm which is almost similar to the context (Joshi and Shrestha, 2019). According to WC measurement, 59.9% (121) of females were abdominally obese while 40.1% (81) had their waist circumference at normal range (below 80 cm). In a cross-sectional study conducted at Kathmandu district reported the prevalence of central obesity among females using Asian criteria for waist circumference was 63.09% which was found higher to its prevalence found in sub-metropolitan city of Itahari (Silvanus, 2018). Figure 4.3 illustrates the prevalence of abdominal obesity by waist circumference.




**Figure 4.3:** Prevalence of obesity with respect to waist circumference

#### 4.6.4 According to waist-to-hip ratio measurement

The prevalence of abdominal obesity was found to be 67.8% (137) while rest 32.2% (65) had normal ratio of less than 0.85. Comparing the results with mean prevalence of 70.2% in Nepal and the mean prevalence of abdominal obesity of 69.4% in province 1 (MOHP *et al.*, 2020), the results was found little lesser in the study. Likewise, with respect to the prevalence of WHR (67.15%) among women in Kathmandu district (Silvanus, 2018), the achieved prevalence was found almost alike but little higher. Another study done in rural districts of Udaypur, south-eastern Nepal, recorded the abdominal obesity of 58% among females (Pyakurel *et al.*, 2019).

Similarly, mean WHR was found to be of  $0.90 \pm 0.06$  cm in the study which was consistent with mean population WHR of 0.90 (MOHP *et al.*, 2020a). A study done in a village at Dolakha, Nepal found the mean WHR among females to be  $0.88 \pm 0.06$  cm (Kharal *et al.*, 2013). Thus, the result was little lower than the average measurement found in sub-metropolitan city of Itahari. Also, a study done in Chitwan also providec little lower mean WHR of  $0.886 \pm 0.054$  cm than the achieved result (Joshi and Shrestha, 2019). Figure 4.4 depicts the prevalence of abdominal obesity by Waist-to-Hip ratio.



**Figure 4.4:** Prevalence of obesity with respect to waist-to-hip ratio

## 4.7 Factors associated with overweight and obesity

### 4.7.1 Factors associated with BMI (WHO cut off)

The chi-square analysis was used to identify the factors associated with overweight and obesity. The analysis showed that age, marital status, parity, sleeping hours, eating pattern/vegetarianism, alcohol, egg and fast foods were significantly associated with BMI. The association is shown in Table 4.14:

The study suggested that the prevalence of overweight and obesity increased as people age. The result of a survey conducted in US concluded that age significantly affect the BMI of adults i.e. increase in age is directly proportional (Canning *et al.*, 2014). Similar result was observed in the study done in Tanzania (George *et al.*, 2021). The association between marital status and obesity can be explained by the fact that people after marriage have less physical activity, change their diet and may be less concerned about their weight. This association was found significantly evident in Greek adults from a national epidemiological survey (Tzotzas *et al.*, 2010). Research done among women living in peri-urban areas in West Africa also highlighted the higher prevalence of overweight and obesity after marriage (Malik *et al.*, 2019).

The study also revealed that overweight or obesity increased with parity. A research based on predictors of overweight and obesity, in Ghana, also explored that having four or more children had a negative impact on the overweight and obesity (Agbeko *et al.*, 2013). Short sleep duration was found significantly associated with increased risk of obesity (M. Watanabe *et al.*, 2010). Short sleep may influence weight gain through different pathways. One mechanism is that short sleep duration leads to increased food intake (Ogilvie and Patel, 2017). Studies of western vegetarians have consistently shown that vegetarians have a lower BMI than otherwise comparable non-vegetarians (Paslakis *et al.*, 2020). However, in an Asian Indian cohort study, it was found that vegetarian dietary patterns were associated with a higher incidence of morbid obesity culminating in bariatric surgery than non-vegetarians (Borude, 2019).

**Table 4.13:** Factors associated with overweight and obesity based on BMI of WHO cut-off (n=202)

<b>Factors</b>	<b>Category</b>	<b>Overweight and obesity frequency</b>	<b>Normal and underweight frequency</b>	<b>P-value</b>
<b>Age</b>	15-20	3 (7.5%)	37 (92.5%)	0.000*
	21-30	23 (34.3%)	44 (65.7%)	
	31-40	24 (47.1%)	27 (52.9%)	
	41-49	37 (84.1%)	7 (15.9%)	
<b>Marital status</b>	Unmarried	6 (10.2%)	53 (89.8%)	0.000*
	Married	80 (56.7%)	61 (43.3%)	
	Widow/ separated	2 (100.0%)	(0.0%)	
<b>Parity</b>	0	14 (18.7%)	61 (81.3%)	0.000*
	1-2	59 (54.6%)	49 (45.4%)	
	≥3	14 (73.7%)	5 .3%)	

<b>Sleep</b>	<7 hours	24 (51.1%)	23 (48.9%)	0.047*
	7-9 hours	54 (39.7%)	82 (60.3%)	
	>9 hours	9 (47.4%)	10 (52.6%)	
<b>Vegetarianism</b>	Vegan	0 (0%)	0 (0%)	0.013*
	Lacto-veg	4 (80.0%)	1 (20.0%)	
	Lacto-ovo-veg	1 (100.0%)	0 (0.0%)	
	Non-veg	82 (41.8%)	114 (58.2%)	
<b>Alcohol</b>	yes	16 (48.5%)	17 (51.5%)	0.087*
	no	62 (36.7%)	107 (63.3%)	
<b>Egg</b>	Regular	14 (43.8%)	18 (56.3%)	0.069*
	Frequent	29 (42.0%)	40 (58.0%)	
	Rare	32 (41.0%)	46 (59.0%)	
	Never	12 (52.2%)	11 (47.8%)	
<b>Fast foods</b>	Regular	21 (38.2%)	34 (61.8%)	0.018*
	Frequent	35 (37.2%)	59 (62.8%)	
	Rare	31 (58.5%)	22 (58.5%)	

\*Statistically significant ( $p < 0.05$ ).

Likewise the increase consumption of fast foods was found to be significantly associated with increased BMI in east Sikkim, India (Kar and Khandelwal, 2015). Alcohol drinking is a component of lifestyle in many regions and, recently, has been raised in some developed countries. It contains high energy (7.1 kcal/g) and also, by disrupting energy balance, leads to accumulation of fat mass and development of overweight or obesity (Golzarand *et al.*, 2021). A community based cross-sectional study in southern Ethiopia defined the association of frequency of egg consumption with overweight or obesity (Darebo *et al.*, 2019).

Other factors such as, occupation, education, monthly income, marital status, types of family, skipping breakfast, eating in front of T.V, meals a day, smoking, consumption fruits and vegetables, polished daal, meats were not significantly associated with prevalence of overweight and obesity.

#### 4.7.2 Factors associated with waist circumference

Age, marital status, education, occupation, family size, parity, contraceptive and physical adequacy were found to have significant association with waist circumference measurement among females in Itahari sub-metropolitan city as shown in Table 4.15.

A population based cohort study among adults found that the incidence of abdominal obesity increases with age as well as dependent on marital status and education level of an individual despite their gender (Barzin *et al.*, 2018). A nationwide survey in Iran highlighted that occupation of females influences the WC in them (Kolahi *et al.*, 2018). Results of the study too signify the relation of WC with occupation of females; in sedentary job holders and women with small business setup like shop-keeping; than in farmers and daily wage workers.

In addition to these factors, parity and contraceptive has also been found to be significant risk factors for central obesity among females. Research done in Jeddah, Saudi among reproductive age women also highlighted the fact with higher WC prevalence in four and more children (Alharbi *et al.*, 2017). In another study among premenopausal women, oral contraceptive has been associated with increased obesity risk (Park *et al.*, 2016). According to study done among the adult population in the Balearic Islands supported the fact of gaining abdominal fat in female after marriage. This could be due to change in dietary patterns, less focus on being attractive, have more social support, being less physically active (Coll *et al.*, 2015).

Different other factors under study; like, alcohol intake, salt intake, monthly income, breakfast skipping, fruits, meat, vegetables and processed food consumption were not found to be significantly associated with abdominal obesity by waist circumference among reproductive aged females.

**Table 4.14:** Factors associated with abdominal obesity of surveyed population (n=202)

<b>Factors</b>	<b>Category</b>	<b>Overweight and obese Frequency</b>	<b>Normal and underweight frequency</b>	<b>P-value</b>
<b>Age</b>	15-20	5 (12.5%)	35 (87.5%)	0.000*
	21-30	38 (78.4%)	29 (43.3%)	
	31-40	40 (78.4%)	11 (21.6%)	
	41-49	38 (86.4%)	6 (13.6%)	
<b>Marital status</b>	Unmarried	10 (16.9%)	49 (83.1%)	0.000*
	Married	109 (77.3%)	32 (22.7%)	
	Widow/ Separated	2 (100%)	0 (0.0%)	
<b>Education</b>	Illiterate	26 (81.2%)	6 (18.8%)	0.001*
	Primary school	23 (79.3%)	6 (20.7%)	
	Secondary school	28 (50%)	28 (50%)	
	High school	33 (54.1%)	28 (45.9%)	
	Graduate	11 (45.8%)	13 (54.2%)	
<b>Occupation</b>	Daily wage worker	13 (92.9%)	1 (7.1%)	0.037*
	Service	10 (66.7%)	5 (33.3%)	
	Farming	7 (70%)	3 (30.0%)	
	Business	11 (68.7%)	5 (31.3%)	
	Unemployed	80 (54.4%)	67 (45.6%)	
<b>Family size</b>	<5	64 (58.7%)	45 (41.3%)	0.018*
	≥5	57 (61.3%)	36 (38.7%)	

<b>Parity</b>	0	21 (28%)	54 (72.0%)	0.000*
	1-2	84 (77.8%)	24 (22.2%)	
	≥3	16 (84.2%)	3 (15.8%)	
<b>Contraceptive</b>	Yes	34 (72.3%)	13 (27.7%)	0.001*
	no	87 (56.1%)	68 (43.9%)	
<b>Physical adequacy</b>	Inadequate	67 (63.2%)	39 (36.8%)	0.047*
	Adequate	54 (56.2%)	42 (43.8%)	

\*Statistically significant (p<0.05)

#### 4.7.3 Factors associated with waist-to-hip ratio

The factors which were significantly associated with waist-to-hip ratio of the participants were age, marital status, education, parity, and contraceptive were associated with WHR.

Abdominal obesity and age was positively associated in the study conducted in Iranian adults (Dalvand *et al.*, 2015). A population based cross-sectional study among Chinese adults also concluded that the prevalence of abdominal obesity increased with age throughout adulthood (Xu *et al.*, 2016). Marital status were found to be associated with WHR in the population based study in China (S. Wu *et al.*, 2014). Likewise, a study conducted in Morocco among child-bearing age women also provided similar results as their level of education been found to be markable determinant factors for increased WHR in them (Barich *et al.*, 2018). Different studies have found significant positive association between central obesity and parity (Owolabi *et al.*, 2017). Also, it was found that both injectable and oral contraceptives were associated with increased WHR among females (Barenson *et al.*, 2009).

Apart from them, factors like family type, watching tv while eating, alcohol intake, GLV and fruit consumption, carbohydrate and fat intake, eating-out of home and consumption of processed food were not found to be statistically significant with abdominal obesity in terms of WHR among survey population.



**Table 4.15:** Factors associated with abdominal obesity based on WHR (n=207)

<b>Factors</b>	<b>Category</b>	<b>Obese frequency (%)</b>	<b>Non-obese frequency (%)</b>	<b>P-value</b>
<b>Age</b>	15-20	8 (20%)	32 (80.0%)	0.000*
	21-30	48 (71.6%)	19 (28.4%)	
	31-40	43 (84.3%)	8 (15.7%)	
	41-49	38 (86.4%)	6 (13.6%)	
<b>Marital status</b>	Unmarried	18 (30.5%)	41 (69.5%)	0.000*
	Married	116 (83%)	24 (17.0%)	
	Widow/ separated	2 (100%)	0 (0.0%)	
<b>Education</b>	Illiterate	28 (87.5%)	4 (12.5%)	0.018*
	Primary school	22 (75.9%)	7 (24.1%)	
	Secondary school	37 (66.1%)	19 (33.9%)	
	High school	33 (54.1%)	28 (45.9%)	
	Graduate	17 (70.8%)	7 (29.2%)	
<b>Parity</b>	0	32 (42.7%)	43 (57.3%)	0.000*
	1-2	89 (82.4%)	19 (17.6%)	
	≥3	16 (84.2%)	3 (15.8%)	
<b>Contraceptive</b>	Yes	37 (78.7%)	10 (21.3%)	0.006*
	no	100 (64.5%)	55 (35.5%)	

\*Statistically significant (p<0.05)

## **Part V**

### **Conclusions and recommendations**

#### **5.1 Conclusions**

The study was intended to identify the risk factors associated with prevalence of overweight and obesity among obesity in 15-49 years male and female residing in Itahari sub-metropolitan city.

- i Factors like age, marital status, parity, sleeping hours, eating pattern/vegetarianism, alcohol, egg and fast foods were associated with BMI. Similarly, Age, marital status, education, occupation, family size, parity, contraceptive and physical adequacy were associated with WC. The factors associated with WHR were age, marital status, education, parity, and contraceptive.
- ii Factors like consumption of alcoholic drinks, sleeping hours, eating pattern/vegetarianism and egg were only associated with BMI. While, physical adequacy and other occupation was only associated with WC.

#### **5.2 Recommendations**

Based on the results obtained in the study, following recommendations could be practiced in order to minimize the risk of overweight and obesity among reproductive aged females residing in Itahari sub- metropolitan city.

1. Physical activity needs to be promoted in females and women-friendly environment needs to be created.
2. Knowledge regarding the portion size and serving sizes should be given so that people will know about their food intake.
3. Further researches can be done in other extended areas of age group, gender specific, and other municipalities and VDC. It can also be extended for other types of schools. Also, comparisons between them can also be done.

## **Part VI**

### **Summary**

A global pandemic of obesity exists, and Nepal is not an exception. The prevalence of overweight and obesity is increasing; and different NCDs are on rise as its consequence. Been evident that women are more obese than men in the whole world, this study focused on obesity prevalence in reproductive age females.

The cross-sectional study was conducted to explore the prevalence and risk factors associated with overweight or obesity among reproductive aged females residing in Itahari sub- metropolitan city. The anthropometric indicators BMI, WC and WHR were used in the study. Along with the socio-demographic, physical activity, behavioral factors, dietary aspects were also explored via structured questionnaires. The data analysis was performed using Excel 2016 and SPSS version 20. Out of 202 respondents, the study reported that 29.2% of respondents were overweight and 9.4% were obese as defined by BMI (WHO criteria). While based on WC and WHR, 59.9% and 67.8% were abdominally obese respectively based on the classification given by IDF and WHO.

The prevalent overweight and obesity in the study site was based on different socio-demographic, economic, behavioral, health and dietary factors under study, age, marital status, parity, sleeping hours, eating pattern/vegetarianism, alcohol, egg and fast foods were significantly associated with BMI. Similarly factors like Age, marital status, education, occupation, family size, parity, contraceptive and physical adequacy were found significantly associated with WC. And finally, the factors associated with WHR were age, marital status, education, parity, and contraceptive.

The reported prevalence of overweight and obesity in Itahari is high and pose a serious health challenge. Thus, timely actions on this issue must be undertaken. Awareness in maintaining healthy body weight, management of associated risk factors and lifestyle intervention programs should be done to combat the prevalent situation.

## References

- Affenito, S. G., Franko, D. L. and Striegel-Moore, R. H. (2012). Behavioral Determinants of Obesity: Research Findings and Policy Implications. *Journal of Obesity*.
- Afshin, A., Forouzanfar, M. H., Sur, P. and Estep, K. (2017). Health effects of overweight and obesity in 195 countries over 25 years. *The New England Journal of Medicine*. **377** (1), 13-27.
- Agbeko, M. P., Akwasi, K.-K. and Andrews, & D. A. (2013). Predictors of Overweight and Obesity Among Women in Ghana. *The Open Obesity Journal*, 72–81.
- Agha, M. and Agha, R. (2017). The rising prevalence of obesity: part A: impact on public health. *International Journal of Surgery, Oncology*. **2** (7), 17.
- Ahmed, M., Seid, A. and Kemal, A. (2020). Does the frequency of watching television matters on overweight and obesity among reproductive age women in Ethiopia? *Journal of Obesity*. **2020**.
- Ahnen, R. T., Jonnalagadda, S. S. and Slavin, J. L. (2019). Role of plant protein in nutrition, wellness, and health. *Nutrition Reviews*. **77** (11), 735-747. 10.1093/nutrit/nuz028.
- Akhil, L. and Ahmad, H. A. (2011). Relationship between Obesity and Cardiovascular Diseases in Four Southern States and Colorado. *Journal of Health Care for Poor and Underserved*. **22** (4), 61-72.
- Albashir, A. A. D. (2020). The potential impacts of obesity on COVID-19. *Clinical medicine*. **20** (4), e109.
- Alberti, G., Zimmet, P., Shaw, J. and Grundy, S. M. (2006). The IDF consensus worldwide definition of the metabolic syndrome. *Brussels: International Diabetes Federation*. **23** (5), 469-480.
- Alharbi, M. and Jackson, & R. T. (2017). Physical Activity and the Prevalence of General and Abdominal Obesity among Saudi Women of Reproductive Age in Jeddah City. *Journal of Obesity and Overweight*, 3, 104.
- Alkerwi, A. a., Crichton, G. E. and Hébert, J. R. (2015). Consumption of ready-made meals and increased risk of obesity: findings from the Observation of Cardiovascular Risk Factors in Luxembourg (ORISCAV-LUX) study. *British Journal of Nutrition*. **113** (2), 270-277.
- Amin, F., Fatima, S. S., Islam, N. and Gilani, A. H. (2015). Prevalence of obesity and overweight, its clinical markers and associated factors in a high risk South-Asian population. *BMC obesity*. **2** (1), 1-11.

- Anderson, B., Lyon-Callo, S., Fussman, C., Imes, G. and Rafferty, A. P. (2011). Peer reviewed: Fast food consumption and obesity among michigan adults. *Preventing chronic disease*. **8** (4).
- Andolfi, C. and Fisichella, P. M. (2018). Epidemiology of obesity and associated comorbidities. *Journal of Laparoendoscopic & Advanced Surgical Techniques*. **28** (8), 919-924.
- Anekwe, C. V., Jarrell, A. R., Townsend, M. J., Gaudier, G. I., Hiserodt, J. M. and Stanford, F. C. (2020). Socioeconomics of Obesity. *Current Obesity Reports*. **9** (3), 272-279. 10.1007/s13679-020-00398-7.
- Aryal, K. K., Neupane, S., Vaidya, A., Singh, S., Paulin, F., Madanlal, R. G., Riley, L. M., Cowan, M., Guthold, R., Singh, S. P., Bhusal, C. L. and Lohani, G. R. (2014). *Non communicable diseases risk factors: STEPS Survey Nepal 2013*. Kathmandu: Nepal Health Research Council
- Ashok, P., Kharche, J. S., Raju, R. and Godbole, G. (2016). Metabolic equivalent task assessment for physical activity in medical students. *National Journal of Physiology, Pharmacy and Pharmacology*. **7** (3), 236-239.
- Baioumi, A. Y. A. A. (2019). Chapter 3 - Comparing Measures of Obesity: Waist Circumference, Waist-Hip, and Waist-Height Ratios. *In: "Nutrition in the Prevention and Treatment of Abdominal Obesity "* (2nd ed.). (R. R. Watson, Ed.). pp. 29-40. Academic Press. [978-0-12-816093-0].
- Barenson, A. B., & Rahman. (n.d.). Changes in weight, total fat, percent body fat, and central-to-peripheral fat ratio associated with injectable and oral contraceptive use. *American Journal of Obstetrics and Gynecology*, 200(329.e321-329.e328).
- Barich, F., Zahrou, F. E, & and Barkat. (2018). Association of Obesity and Socioeconomic Status among Women of Childbearing Age Living in Urban Area of Morocco. *Journal of Nutrition and Metabolism*.
- Baker, A., Sirois-Leclerc, H. and Tulloch, H. (2016). The Impact of Long-Term Physical Activity Interventions for Overweight/Obese Postmenopausal Women on Adiposity Indicators, Physical Capacity , and Mental health Outcomes: A Systematic Review. *Journal of Obesity*. 22.
- Barzin, M., Piri, Z., Serahati, S., Valizadeh, M., Azizi, F. and Hosseinpanah, F. (2018). Incidence of abdominal obesity and its risk factors among Tehranian adults. *Public health nutrition*. **21** (17), 3111-3117.
- Beccuti, G. and Pannain, S. (2011). Sleep and obesity. *Current opinion in clinical nutrition and metabolic care*. **14** (4), 402.
- Bechthold, A. (2014). Food energy density and body weight. *Ernahrungs Umschau*. **6**, 2-11.

- Bendsen, N. T., Christensen, R., Bartels, E. M., Kok, F. J., Sierksma, A., Raben, A. and Astrup, A. (2013). Is beer consumption related to measures of abdominal and general obesity? A systematic review and meta-analysis. *Nutrition reviews*. **71** (2), 67-87.
- Bezerra, I., Curioni, C. and Sichieri, R. (2012). Association between eating out of home and body weight. *Nutrition Reviews*. **70** (2), 65-79.
- Bhandari, S., Sayami, J. T., Thapa, & P. and Sayami. (2016). Dietary intake patterns and nutritional status of women of reproductive age in Nepal: findings from a health survey. *Archives of Public Health*, **7**
- Bhurosy, T. and Jeewon, R. (2014). Overweight and obesity epidemic in developing countries: a problem with diet, physical activity, or socioeconomic status? *The Scientific World Journal*, 2014.
- Bhutani, S., Schoellar, D. A. and Walsh, M. C. (2016). Frequency of eating out at both fast-food and sit-down restaurants was associated with high body mass index in non-large metropolitan communities in Midwest. *American Journal of Health Promotion*. **32** (1), 75-83.
- Blair, S. N. and Moriss, J. N. (2009). Healthy hearts and the universal benefits of being physically active: physical activity and health. *Annals of Epidemiology*. **19** (4), 253-256.
- Bonanno, L., Metro, D., Papa, M., Finzi, G., Maviglia, A., Sottile, F., Corallo, F. and Manasseri, L. (2019). Assessment of sleep and obesity in adults and children: Observational study. *Medicine*. **98** (46).
- Borude, & S. (2019). Which Is a Good Diet—Veg or Non-veg? Faith-Based Vegetarianism for Protection From Obesity—a Myth or Actuality? *Obesity Surgery*, **29**, 1276–1280.
- Bray, G. A. and Bouchard, C. (2019). "Handbook of Obesity" (Fourth ed.). CRC Press.
- Bryan, S. and Walsh, P. (2004). Physical Activity and Obesity in Canadian Women. *BMC Women's Health*. **4** (1), 1-10.
- Bujnowski, D., Xun, P., Daviglius, M. L., Van Horn, L., He, K. and Stamler, J. (2011). Longitudinal Association between Animal and Vegetable Protein Intake and Obesity among Men in the United States: The Chicago Western Electric Study. *Journal of the American Dietetic Association*. **111** (8), 1150-1155.e1151. <https://doi.org/10.1016/j.jada.2011.05.002>.
- Byeong, Park, H. A., Kang, J.-H., Kim, K., Cho, Y. G. and Jang, J. (2018). Animal and plant protein intake and body mass index and waist circumference in a Korean elderly population. **10** (5), 577.

- Canning, K. L., Brown, R. E., Jamnik, V. K. and Kuk, J. L. (2014). Relationship between obesity and obesity-related morbidities weakens with aging. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*. **69** (1), 87-92.
- Carmen, S.-O., Miguel A, M. G. and Maira, B. R. (2011). Alcohol consumption and body weight: a systematic review. *Nutrition reviews*. **69** (8), 419-431.
- CBS. (2021). National Census 2021 [Report]. Central Bureau of Statistics. Kathmandu, Nepal.
- Chaput, J.-P., Klingenberg, L., Rosenkilde, M., Gilbert, J.-A., Tremblay, A. and Sjödín, A. (2011). Physical Activity Plays an Important Role in Body Weight Regulation. *Journal of Obesity*. 10.1155/2011/360257.
- Chaudhary, S. (2020). RISK FACTORS ASSOCIATED WITH OVERWEIGHT AND OBESITY AMONG 18-59 YEARS MALE AND FEMALE RESIDING IN RAMDHUNI MUNICIPLITY, SUNSARI. Department Of Nutrition and Dietetics Central Campus of Technology ...,
- Choi, J., Guterrez, Y., Gilliss, C. and Lee, K. A. (2012). Physical activity, weight, and waist circumference in midlife women. *Health Care for Women International*. **33** (12), 1086-1095.
- Chowdhury, E. A., Richardson, J. D., Holman, G. D., Tsintzas, K., Thompson, D. and Betts, J. A. (2016). The causal role of breakfast in energy balance and health: a randomized controlled trial in obese adults. *The American Journal of Clinical Nutrition*. **103** (3), 747-756. 10.3945/ajcn.115.122044.
- Coll, J. L., Bibiloni Mdel, M., Salas, Pons, A. and Tur, & J. A. (2015). *Prevalence and Related Risk Factors of Overweight and Obesity among the Adult Population in the Balearic Islands, a Mediterranean Region*. 220–233.
- Cooper, C. B., Neufeld, E. V., Dolezal, B. A. and Martin, J. L. (2018). Sleep deprivation and obesity in adults: a brief narrative review. *BMJ open sport & exercise medicine*. **4** (1), e000392.
- Coronel, J., Pinos, I. and Amengual, J. (2019).  $\beta$ -carotene in obesity research: Technical considerations and current status of the field. *Nutrients*. **11** (4), 842.
- Csige, I., Ujvarosy, D. and Szabo, Z. (2018). The Impact of Obesity on the Cardiovascular System. *Journal of Diabetes Research*.
- Dabrowska, J., Dabrowska-Galas, M., Naworska, B. and Woderska, M. (2015). The role of physical activity in preventing obesity in midlife women. *Prezegland Menopauzalny*. **14** (1), 13-19.

- Dalvand, S., Koohpayehzadeh, J., Karimlou, M., Asgari, F., Rafei, A., Seifi, B., Niksima, S. H. and Bakhshi, E. (2015). Assessing factors related to waist circumference and obesity: Application of a latent variable model. *Journal of environmental and public health*. **2015**.
- Darebo, T., Mesfin, A. and Gabremedhin, & S. (2019). Prevalence and factors associated with overweight and obesity among adults in Hawassa city, southern Ethiopia: a community based cross-sectional study. *BMC Obesity*.
- Depres, J. P. and Lemieux, I. (2006). Abdominal obesity and metabolic syndrome. *Nature*. 444:881-887.
- DFTQC. (2017). "Food Composition Table For Nepal" (Third ed.). Department of Food Technology and Quality Control. Babarmahal, Kathmandu.
- Dhimal, M., Bista, B., Bhattarai, S., Dixit, L. P., Hyder, M. K. A., Agrawal, N., Rani, M. and Jha, A. K. (2020). Report of Noncommunicable Disease Risk Factors: STEPS Survey Nepal 2019. Kathmandu: Nepal Health Research Council.
- Dinour, L., Leung, M. M., Tripicchio, G., Khan, S. and Yeh, M.-C. (2012). The association between marital transitions, body mass index, and weight: a review of the literature. *Journal of obesity*. **2012**.
- Doo, M. and Kim, Y. (2017). The risk of being obese according to short sleep duration is modulated after menopause in Korean women. **9** (3), 206.
- Drewnowski, A. (2007). The real contribution of added sugars and fats to obesity. *Epidemiologic reviews*. **29** (1), 160-171.
- Drewnowski, A. (2009). Obesity, diets, and social inequalities. *Nutrition reviews*. **67** (suppl\_1), S36-S39.
- Fazzino, T. L., Fleming, K., Sher, K. J., Sullivan, D. K. and Befort, C. (2017). Heavy drinking in young adulthood increases risk of transitioning to obesity. *American journal of preventive medicine*. **53** (2), 169-175.
- Filgueiras, M. D., Vieira, S. A. and Fonseca, P. C. D. A. (2019). Waist circumference, waist-to-height ratio and conicity index to evaluate android fat excess in Brazilian children. *Public Health Nutrition*. **22** (1), 140-146.
- Gakidou, E. (2014). Prevalence of overweight and obesity in children and adults—Authors' reply. **384** (9960), 2108.



- Gallagher, D., Heymsfield, S. B., Heo, M., Jebb, S. A., Murgatroyd, P. R. and Sakamoto, Y. (2000). Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *The American journal of clinical nutrition*. **72** (3), 694-701.
- Gandy, W. J., Madden, A. and Holdsworth, M. (2006). "Oxford Handbook of Nutrition and Dietetics" (1st ed.). Oxford University Press. New York. [978-0-19-856725-7].
- George, J. M., Mpogole, Z., Mgongo, M., Mamseri, R., Leyaro, B. J., Mauka, W. and Msuya, S. E. (2021). Prevalence and Factors Associated With Overweight or Obesity Among Primary and Secondary School Teachers In Moshi Municipality, Kilimanjaro, Tanzania.
- Ghosh, A. and Ghosh, T. (2009). Modification of Kuppuswamy's Socioeconomic Status Scale in context to Nepal.
- Gluckman, P., Nishtar, S. and Armstrong, T. (2015). Ending childhood obesity: a multidimensional challenge. *The Lancet*. **385** (9973), 1048-1050.
- Golay and Bobbioni, E. (1997). The role of dietary fat in obesity. **21**, S2-11.
- Golzarand, M., Salari-Moghaddam, A. and Mirmiran, P. (2021). Association between alcohol intake and overweight and obesity: a systematic review and dose-response meta-analysis of 127 observational studies. *Critical Reviews in Food Science and Nutrition*. 1-21.
- Gupta, P., Tyagi, S. and Mukhiya, M. (2011). Obesity: An Introduction and Evaluation. *Journal of American College of Nutrition*. **21** (2), 152-155.
- Hales, C. M., Fryar, C. D., Carroll, M. D., Freedman, D. S. and Ogden, C. L. (2018). Trends in obesity and severe obesity prevalence in US youth and adults by sex and age, 2007-2008 to 2015-2016. *Jama*. **319** (16), 1723-1725.
- Helble, M. and Francisco, K. (2017). The upcoming obesity crisis in Asia and the Pacific: First cost estimates. *Tokyo: Asian Development Bank Institute: ADBI Working Paper*. **743**.
- Hill, J. O. and Peters, J. C. (1998). Environmental contributions to the obesity epidemic. *Science*. **280** (5368), 1371-1374.
- Hopkin, M. (2008). Fat cell numbers stay constant through adult life. *Nature*. 10.1038/news.2008.800.
- Hwang, Y., Hayashi, T., Fujimoto, W., Kahn, S., Leonetti, D., McNeely, M. and Boyko, E. (2015). Visceral abdominal fat accumulation predicts the conversion of metabolically healthy obese subjects to an unhealthy phenotype. *International journal of obesity*. **39** (9), 1365-1370.

- ICMR. (2010). Nutrient Requirements and Recommended Dietary Allowances for Indians [Report]. National Institution of Nutrition. Hyderabad India.
- IDF. (2006). The IDF consensus worldwide definition of the Metabolic Syndrome [Report]. International Diabetes Federation. Brussels, Belgium,
- IDF.2005. The IDF consensus worldwide definition of the metabolic syndrome. Retrieved from [www.idf.org.communicationidf.org](http://www.idf.org.communicationidf.org). (Last update 2005). [Accessed 25 may,2022].
- IPAQ. (2005). Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ). In IPAQ (Ed.). (Revised ed., pp. 15 pages).
- Jacqui, W. (2017). Waist measurement, not BMI, is stronger predictor of death risk, study finds. *British Medical Journal*. **357**.
- James, W. (2008). The epidemiology of obesity: the size of the problem. *Journal of Internal Medicine*. **263** (4), 336-352.
- Janjua et al., Mahmood, & Bhatti. (2015). ). Association of Household and Community Socioeconomic Position and Urbanicity with Underweight and Overweight among Women in Pakistan. *PLoS ONE*, *10*.
- Jensen, M. D., Ryan, D. H., Apovian, C. M., Ard, J. D., Comuzzie, A. G., Donato, K. A., Hu, F. B., Hubbard, V. S., Jakicic, J. M. and Kushner, R. F. (2014). 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Journal of the American college of cardiology*. **63** (25 Part B), 2985-3023.
- Jiménez, E. G. (2013). Obesity: etiologic and pathophysiological analysis. *Endocrinología y Nutrición (English Edition)*. **60** (1), 17-24.
- Joshi, B. and S. L. (2019). A Comparative Study of Waist Hip Ratio and Body Mass Index (BMI) in Diabetic and Non Diabetic Individuals of Chitwan, Nepal. *Journal of Diabetes & Metabolism*.
- Joshi, B., Shrestha, L., Bhattarai, K., Manandhar, N. and Mahotra, N. B. (2019). Comparison of central obesity with overall obesity in predicting the risk of type 2 diabetes mellitus. *Journal of Universal College of Medical Sciences*. **7** (1), 17-21.
- Jura, M. and Kozak, L. P. (2016). Obesity and related consequences to ageing. *Age (Dordr)*. **38** (1), 23.

- Kar, S. and Khandelwal, B. (2015). Fast foods and physical inactivity are risk factors for obesity and hypertension among adolescent school children in east district of Sikkim, India. *Journal of natural science, biology, and medicine*. **6** (2), 356.
- Kharal, P. M., Prasad, & P. N. and Acharya. (2013). Gross Correlation between Waist Hip Ratio and Blood Sugar Level in a Village. *Journal of Nepal Medical Association.*, **52**, 361–364.
- Kibria, G. M. A. (2019). Prevalence and factors affecting underweight, overweight and obesity using Asian and World Health Organization cutoffs among adults in Nepal: Analysis of the Demographic and Health Survey 2016. *Obesity Research & Clinical Practice*, **13**(2), 129–136.
- Kikuchi, Y., Nozaki, S., Makita, M., Yokozuka, S., Fukudome, S.-i., Yanagisawa, T. and Aoe, S. (2018). Effects of whole grain wheat bread on visceral fat obesity in Japanese subjects: A randomized double-blind study. *Plant Foods for Human Nutrition*. **73** (3), 161-165.
- Koirala, M. , B. S. and K. M. L. (2019). Risk factors for Obesity in Nepalese Women: A Cross-sectional Study. *Journal of Lumbini Medical College*, **7**, 93–99.
- Kolahi, A.-A., Moghisi, A. and Ekhtiari, & Y. (2018). Socio-demographic determinants of obesity indexes in Iran: findings from a nationwide STEPS survey. *Health Promotion Perspectives*, **8**(3), 187–194.
- Kranendonk, M. E. G., van Herwaarden, J. A., Stupkova, T., Jager, W. d., Vink, A., Moll, F. L., Kalkhoven, E. and Visseren, F. L. J. (2015). Inflammatory characteristics of distinct abdominal adipose tissue depots relate differently to metabolic risk factors for cardiovascular disease: Distinct fat depots and vascular risk factors. *Atherosclerosis*. **239** (2), 419-427. <https://doi.org/10.1016/j.atherosclerosis.2015.01.035>.
- Kumar, A., Kumar, P. and Joshi, P. (2016). Food consumption pattern and dietary diversity in Nepal: implications for nutrition security. *Indian Journal of Human Development*. **10** (3), 397-413.
- Kumar, & P. and Joshi. (2016). Food consumption pattern and dietary diversity in Nepal: implications for nutrition security. *Indian Journal of Human Development*, **10**, 397–413.
- Kumar et al. (2016). *Food consumption pattern and dietary diversity in Nepal: implications for nutrition security*.
- Kushner, R. F. (2012). Clinical Assessment and Management of Adult Obesity. *Circulation*. **126**, 2870-2877.

- Kushner, R. F. and Ryan, D. (2014). Assessment and lifestyle management of patients with obesity: clinical recommendations from systematic reviews. *JAMA*. **312**, 943-952.
- Larsen, S. C., Ängquist, L., Sørensen, T. I. and Heitmann, B. L. (2013). 24h urinary sodium excretion and subsequent change in weight, waist circumference and body composition. *PloS one*. **8** (7), e69689.
- Lee, A., Cardel, M. and Donahoo, W. T. (2000). "Social and Environmental Factors Influencing Obesity". MDText.com, Inc., South Dartmouth (MA).
- Leung, M. Y. M., Carlsson, N. and Colditz, G. A. (2017). The Burden of Obesity on Diabetes in the United States: Medical Expenditure Panel Survey , 2008-2012. *Value Health*. **20** (1), 77-84.
- Liu, W., Zhang, R., Tan, A., Ye, B., Zhang, X., Wang, Y., Zou, Y., Ma, L., Chen, G. and Li, R. (2019). Long sleep duration predicts a higher risk of obesity in adults: a meta-analysis of prospective cohort studies. *Journal of Public Health*. **41** (2), e158-e168.
- López-Sobaler, A. M., Rodríguez-Rodríguez, E., Aranceta-Bartrina, J., Gil, Á., González-Gross, M., Serra-Majem, L., Varela-Moreiras, G. and Ortega, R. M. (2016). General and abdominal obesity is related to physical activity, smoking and sleeping behaviours and mediated by the educational level: findings from the ANIBES study in Spain. **11** (12), e0169027.
- Malik, K. S., & Adoubi, A. and K. (2019). Overweight and obesity among women living in peri-urban areas in West Africa. *South Eastern European Journal of Public Health*, **12**.
- Ma, W.-Y., Yang, C.-Y., Shih, S.-R., Hsieh, H.-J., Hung, C. S., Chiu, F.-C., Lin, M.-S., Liu, P.-H., Hua, C.-H., Hsein, Y.-C., Chuang, L.-M., Lin, J.-W., Wei, J.-N. and Li, H.-Y. (2013). Measurement of Waist Circumference: Midabdominal or iliac crest? *Diabetes Care*. **36** (6), 1660-1666. 10.2337/dc12-1452.
- Ma, X., Chen, Q., Pu, Y., Guo, M., Jiang, Z., Huang, W., Long, Y. and Xu, Y. (2020). Skipping breakfast is associated with overweight and obesity: A systematic review and meta-analysis. *Obesity Research & Clinical Practice*. **14** (1), 1-8. <https://doi.org/10.1016/j.orcp.2019.12.002>.
- Ma, Y., He, F. J. and MacGregor, G. A. (2015). High salt intake: independent risk factor for obesity? *Hypertension*. **66** (4), 843-849.
- Mahan, L. K. and Raymond, J. L. (2016). "Krause's food & the nutrition care process-e-book". Elsevier Health Sciences. [0323340768].
- Mahan, L. K. and Raymond, J. L. (2017). "Krause's Food and The Nutrition Care Process" (14th ed.). Elsevier Inc. St. Louis, Missouri.

- Mazzocoli, G. (2016). Body composition: Where and When. *European Journal of Radiology*. **85** (8), 1456-1460.
- McArdle, W. D., Katch, F. I. and Katch, V. L. (2010). "Exercise Physiology: Nutrition, Energy, and Human Performance" (Seventh ed.). Lippincott Williams & Wilkins. Walnut Street, Philadelphia. [0781797810].
- McKeown, N. M., Yoshida, M. and Shea, M. K. (2009). Whole Grain Intake and Cereal Fiber are associated with Lower Abdominal Adiposity in Older Adults. *The Journal of Nutrition*. **139** (10), 1950-1955.
- McMillen, M. (2017). The Connection Between Salt and Weight. Web. MD.
- Mendoza, J. A., Drewnowski, A. and Christakis, D. A. (2007). Dietary energy density is associated with obesity and the metabolic syndrome in US adults. *Diabetes care*. **30** (4), 974-979.
- Merchant, Vatanparast, H., Barlas, S., Dehghan, M., Shah, S. M. A., De Koning, L. and Steck, S. E. (2009). Carbohydrate intake and overweight and obesity among healthy adults. **109** (7), 1165-1172.
- Merritt, D. C., Jamnik, J. and El-Sohemy, A. (2018). FTO genotype, dietary protein intake, and body weight in a multiethnic population of young adults: a cross-sectional study. *Genes & nutrition*. **13** (1), 1-10.
- Mohammadbeigi, A., Asgarian, A., Moshir, E., Heidari, H., Afrashteh, S., Khazaei, S. and Ansari, H. (2018). Fast food consumption and overweight/obesity prevalence in students and its association with general and abdominal obesity. *Journal of preventive medicine and hygiene*. **59** (3), 236.
- MOHP and ICF. (2017). *National Demographic and Health Survey, 2016*. Kathmandu, Nepal: Ministry of Health, Nepal.
- MOHP, Nepal, G. o., SOLID Nepal and WHO. (2009). WHO STEPS Surveillance: Non Communicable Disease Risk Factors Survey, 2008 [Report].
- MOPH. (2013). Non communicable diseases risk factors: steps survey Nepal. 2013.
- MOHP, N. and W. (2020b). *Noncommunicable Disease Risk Factors: STEPS Survey Nepal 2019*.
- Mozaffarian, D., Hao, T., Rimm, E. B., Willett, W. C. and Hu, F. B. (2011). Changes in diet and lifestyle and long-term weight gain in women and men. *New England Journal of Medicine*. **364** (25), 2392-2404.

- Municipality, I. (2019). Itahari sub-metropolitan city office Itahari Sub-metropolitan City. Retrieved from <https://itaharimun.gov.np/ne/content>.
- Must, A., McKeown, N. M., Groot, L. J., Beck-Peccoz, P., Chrousos, G. and Dungan, K. (2012). The disease burden associated with overweight and obesity. *Endotext. MDText.com, Inc.; 2000-2012, South Dartmouth, MA.*
- National Planning Commission, N. and Central Bureau of Statistics, C. (2013). Nepal Thematic Report on Food Security and Nutrition 2013. NPC and CBS. Kathmandu, Nepal,
- NCD-RisC. (2016). Trends in adult body-mass in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 participants. *The Lancet.* **387** (10026), 1377-1396.
- Nepal Rastra Bank, N. (2016). *Fifth Household Budget Survey Nepal 2014/15.*
- OECD and WHO. (2020). "Overweight and obesity".
- Ogilvie, R. P. and Patel, S. R. (2017). The epidemiology of sleep and obesity. *Sleep Health.* **3** (5), 383-388.
- Oh, S. W., Han, K. H., Han, S. Y., Koo, H. S., Kim, S. and Chin, H. J. (2015). Association of sodium excretion with metabolic syndrome, insulin resistance, and body fat. *Medicine.* **94** (39).
- Oh, S. W., Koo, H. S., Han, K. H., Han, S. Y. and Chin, H. J. (2017). Associations of sodium intake with obesity, metabolic disorder, and albuminuria according to age. *PloS one.* **12** (12), e0188770.
- Ohno, Y., Hirai, K., Sato, N. and Ito, M. (1997). Food consumption pattern and nutrient intake among Nepalese living in the southern rural Terai region. *Asia Pacific Journal of Clinical Nutrition.* **6** (4), 251-255.
- Okorodudu, D. O., Jumean, M. F., Montori, V. M. and Corral, A. R. (2010). Diagnostic Performance of Body Mass Index to Identify Obesity as Defined by Body Adiposity: A Systematic Review and Meta-Analysis. *International Journal of Obesity.* **34** (5), 791-799.
- Önnerfält, J., Erlanson-Albertsson, C., Montelius, C. and Thorngren-Jerneck, K. (2018). Obese children aged 4–6 displayed decreased fasting and postprandial ghrelin levels in response to a test meal. *Acta Paediatrica.* **107** (3), 523-528.
- Ouchi, N., Parker, J. L., Lugus, J. J. and Walsh, K. (2011). Adipokines in inflammation and metabolic disease. *Nature reviews immunology.* **11** (2), 85-97.

- Owolabi, E. O., Goon, D. T. and Adeniyi, & O. V. (2017). Central obesity and normal-weight central obesity among adults attending healthcare facilities in Buffalo City Metropolitan Municipality, South Africa: a cross-sectional study. *Journal of Health, Population and Nutrition*, 36.
- Öztürk, Z. A., Türkbeyler, İ. H., Abiyev, A., Kul, S., Edizer, B., Yakaryılmaz, F. D. and Soylu, G. (2018). Health-related quality of life and fall risk associated with age-related body composition changes; sarcopenia, obesity and sarcopenic obesity. *Internal medicine journal*. **48** (8), 973-981.
- Park, & B. and Kim. (2016). Oral Contraceptive Use, Micronutrient Deficiency, and Obesity among Premenopausal Females in Korea: The Necessity of Dietary Supplements and Food Intake Improvement. *PLoS ONE*, 11(e0158177).
- Paslakis, G., Richardson, C. and Nohre, & M. (2020). *Prevalence and psychopathology of vegetarians and vegans – Results from a representative survey in Germany*.
- Patil, Y. N., Dandekar, V. and Padghan, P. (2021). Role of Milk and Milk Products in Human Health. *Food and Scientific reports*. **2** (1), 28.
- Piryani, Suneel, Baral, K. P., Pradhan, B., Poudyal, A. K. and Piryani, R. M. (2016a). Overweight and its associated risk factors among urban school adolescents in Nepal: a cross-sectional study. **6** (5), e010335.
- Piryani, S., Baral, K., Pradhan, B. and Poudyal, A. (2016b). Overweight and its associated risk factors among urban school adolescents in Nepal: A Cross-Sectional Study. *BMJ Open*. **6** (5).
- Popkin, B. M., Adair, L. S. and Ng, S. W. (2012). Global nutrition transition and the pandemic of obesity in developing countries. *Nutrition reviews*. **70** (1), 3-21.
- Prentice, A. M. and Jebb, S. A. (2003). Fast foods, energy density and obesity: a possible mechanistic link. *Obesity reviews*. **4** (4), 187-194.
- Purnell, J. Q. (2018). Definitions, Classification, and Epidemiology of Obesity. *MDText.com, Inc: South Dartmouth*.
- Pyakurel, P., Yadav, D. K. and Thapa, J. (2019). Prevalence and associated risk factor of hypertension among individuals of age 18-59 years in South-eastern Nepal: A cross-sectional study. *Nepalese Heart Journal*. **16** (1), 19-26.
- Ramirez-Velez, R., Correa-Bautista, J. E. and Sanders-Tordecilla, A. (2017). Percentage of Body Fat and Fat Mass Index as a Screening Tool for Metabolic Syndrome Prediction in Colombian University Students. *Nutrients*. **9** (9), 1009.

- Rautiainen, S., Wang, L., Lee, I.-M., Manson, J. E., Buring, J. E. and Sesso, H. D. (2016). Dairy consumption in association with weight change and risk of becoming overweight or obese in middle-aged and older women: a prospective cohort study. *The American journal of clinical nutrition*. **103** (4), 979-988.
- Rawal, L. B. , K. K. and J. D. (2018). *Prevalence of underweight, overweight and obesity and their associated risk factors in Nepalese adults: Data from a Nationwide Survey*. *13*(e205912).
- Ritchie, H. and Roser, M. (2020). Obesity. Our world in data.
- Rollins, K., Awwad, A. and Macdonald, I. (2017). Body composition measurement using computed tomography: Does the phase scan matter? *Nutrition*. **41** (37-44). 10.1016/j.nut.2017.02.011.
- Rolls, B. J., Ello-Martin, J. A. and Tohill, B. C. (2004). What can intervention studies tell us about the relationship between fruit and vegetable consumption and weight management? *Nutrition reviews*. **62** (1), 1-17.
- Ronti, T., Lupattelli, G. and Mannarino, E. (2006). The endocrine function of adipose tissue: an update. *Clin Endocrinol*. *64*:355-365.
- Rovira, R. F. (2015). Milk and milk products: food sources of calcium. *Nutricion hospitalaria*. **31**, 1-9.
- Sahakyan, K. R., Somers, V. K., Rodriguez-Escudero, J. P., Hodge, D. O., Carter, R. E., Sochor, O., Coutinho, T., Jensen, M. D., Roger, V. L. and Singh, P. (2015). Normal-weight central obesity: implications for total and cardiovascular mortality. **163** (11), 827-835.
- Sahoo, K., Sahoo, B., Choudhary, A. and Sofi, N. Y. (2015). Childhood obesity: causes and consequences. *Journal of Family Medicine and Primary Care*. **4** (2), 187-192.
- Sangachin, M. G., Cavuto, L. A. and Wang, Y. (2018). Use of various obesity measurement and classification methods in occupational safety and health research: a systematic review of the literature. *BMC Obesity*. **5** (28).
- Sartorius, B., Veerman, L. J., Manyema, M., Chola, L. and Hofman, K. (2015). Determinants of obesity and associated population attributability, South Africa: Empirical evidence from a national panel survey, 2008-2012. *PloS one*. **10** (6), e0130218.
- Sato, A. P. S., Fujimori, E., Szarfarc, S. C., Borges, A. L. V. and Tsunehiro, M. A. (2010). Food consumption and iron intake of pregnant and reproductive aged women. **18**, 247-254.



- Seguin, R. A., Aggarwal, A. and Vermeulen, F. (2016). Consumption Frequency of Foods Away from Home Linked with Higher Body Mass Index and Lower Fruit and Vegetable Intake among Adults: A Cross-Sectional Study. *Journal of Environmental and Public Health*.
- Seidell, J. (2010). Waist Circumference and waist/hip ratio in relation to all-cause mortality, cancer and sleep apnea. *European Journal of Clinical Nutrition*. **64** (1), 35-41.
- Shahi, M., Rai, L., Adhikari, R. D. and Sharma, M. (2013a). Prevalence and factors associated with obesity among adult women of Nepal. *Global Journal of Medicine and Public, GJMEDPH*. **2** (4), 1-9.
- Shahi, M., Rai, L., Adhikari, R. D. and Sharma, M. (2013b). Prevalence and factors associated with obesity among adult women of Nepal. **2** (4), 1-9.
- Sharma, Kumar, S., Ghimire, A., Radhakrishnan, J., Thapa, L., Shrestha, N. R., Paudel, N., Gurung, K., Budathoki, A., Baral, N. and Brodie, D. (2011a). Prevalence of hypertension, obesity, diabetes, and metabolic syndrome in Nepal. **2011**.
- Sharma, S. K., Ghimire, A., Radhakrishnan, J., Thapa, L., Shrestha, N. R., Paudel, N., Gurung, K., Budathoki, A., Baral, N. and Brodie, D. (2011b). Prevalence of hypertension, obesity, diabetes, and metabolic syndrome in Nepal. **2011**.
- Sheth, D. M. and Shah, M. N. (2006). "The Scientific Way to Managing Obesity". Sterling Publishers(P) Ltd. New Delhi.
- Shrestha, S. (2017). Assessment of behavioral risk factors of non-communicable diseases among higher secondary school teachers of Kathmandu. *International Journal of Health Sciences and Research*. **7** (3), 269-273.
- Silvanus, V., Gupta, R. and Dhakal, N. (2018). Comparison of anthropometric indices as correlates of obesity and hypertension among an adult population in the Kathmandu district, Nepal. *Nepal Med Coll J*. **20** (1-3), 52-67.
- Simkhada, P., Poobalan, A., Simkhada, P. P., Amalraj, R. and Aucott, L. (2011). Knowledge, attitude, and prevalence of overweight and obesity among civil servants in Nepal. *Asia Pacific Journal of Public Health*. **23** (4), 507-517.
- Slavin, J. L. and Lloyd, B. (2012). Health Benefits of Fruits and Vegetables. *Advances in Nutrition*. **3** (4), 506-516. 10.3945/an.112.002154.
- Smolnikova, F., Rebezov, M., Chernenkov, E., Sharipova, A., Zhilyaev, R., Anichkina, O., Babaeva, M., Voskanyan, O. and Vorobyev, D. (2021). Developing milk-fruit drinks for school nutrition. *Journal of Natural Remedies*. **21** (9 (1)), 72-77.

- Spalding, K. L., Arner, E., Westermark, P. O. and Bernard, S. (2008). Dynamics of fat cell turnover in humans. *Nature*. **453** (7196), 783-787.
- Srilakshmi. (2019). "Dietetics" (Eighth ed.). New Age International (P) Limited. Daryaganj, New Delhi.
- Stone, T. W., McPherson, M. and Darlington, L. G. (2018). Obesity and Cancer: Existing and New Hypothesis for a Casual Connection. *EBio Medicine*. **30**, 14-28.
- Subedi, Y. P., Marais, D. and Newlands, D. (2017). Where is Nepal in the nutrition transition? *Asia Pacific journal of clinical nutrition*. **26** (2), 358-367.
- Teachman, J. (2016). Body weight, marital status, and changes in marital status. *Journal of family issues*. **37** (1), 74-96.
- Thorp, A., Owen, N. and Neuhas, M. (2011). Sedentary behaviors and subsequent health outcomes in adults : A systematic review of longitudinal studies 1996-2011. *American Journal of Preventive Medicine*. **41** (2), 207-215.
- Tilg, H. and Moschen, A. (2006). Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nature Reviews, Immunology*. **6** (10), 772-783.
- Trang, L. T., Trung, N. N. and Chu, D. T. (2019). Percentage Body Fat as a Good Indicator for Determining Adolescents who are Overweight and Obese: A cross-sectional study in Vietnam. *Osong Public Health and Reasearch Perspectives*. **10** (2), 108-114.
- Traversy, G. and Chaput, J.-P. (2015). Alcohol consumption and obesity: an update. *Current obesity reports*. **4** (1), 122-130.
- Tzotzas, T., Vlahavas, & G. and Papadopoulou. (2010). Marital status and educational level associated to obesity in Greek adults: data from the National Epidemiological Survey. *BMC Public Health*., **10**.
- Umuerri, E. M. (2019). Chapter 17 - Ethnicity and Cut-Off Values in Obesity. *In: "Nutrition in the Prevention and Treatment of Abdominal Obesity "* (2nd ed.). (R. R. Watson, Ed.). pp. 211-223. Academic Press. [978-0-12-816093-0].
- Vaidya, Abhinav, Shakya, S. and Krettek, A. (2010). Obesity prevalence in Nepal: public health challenges in a low-income nation during an alarming worldwide trend. **7**
- Vincente, A. R., Manganaris, G. A., Ortiz, C. M., Sozzi, G. O. and Crisosto, C. H. (2014). Chapter 5 - Nutritional Quality of Fruits and Vegetables. *In: "Postharvest Handling (Third Edition)"* (3rd ed.). (W. J. Florkowski, R. L. Shewfelt, B. Brueckner and S. E. Prussia, Eds.). pp. 69-122. San Diego. Academic Press. [978-0-12-408137-6].

- Watanabe, M., Kikuchi, H., Tanaka, K. and Takahashi, M. (2010). Association of short sleep duration with weight gain and obesity at 1-year follow-up: a large-scale prospective study. *Sleep*. **33** (2), 161-167.
- Watanabe, Y., Saito, I., Henmi, I., Yoshimura, K., Maruyama, H., Yamauchi, K., Matsuo, T., Kato, T., Tanigawa, T. and Kishida, T. (2014). Skipping breakfast is correlated with obesity. *Journal of Rural Medicine*. 2887.
- Watson, S. and Bubnis, D. (2017). What is the Waist-to-Hip Ratio? *Healthline*.
- Whitney, E. and Rolfes, R. S. (2011). "Understanding Nutrition" (12th ed.). Wadsworth Cengage Learning. Belmont, USA. [978-0-538-49412-0].
- WHO. (2003). Global Strategy on Diet, Physical Activity and Health [Report]. Geneva, Switzerland.
- WHO. (2004). WHO global strategy on diet, physical activity and health. **25** (3), 292-302.
- WHO. (2005). WHO STEPS surveillance manual: the WHO STEPwise approach to chronic disease risk factor surveillance [Report]. 9241593830. World Health Organization. Geneva,
- WHO. (2010). Global recommendations on physical activity for health [Report]. Geneva,
- WHO. (2011). Waist circumference and waist-hip ratio: report of a WHO expert consultation [Report]. Geneva,
- WHO. (2012). Guideline: Sodium intake for adults and children [Report]. Geneva, Switzerland.
- WHO. (2013). Guidance on emerging double threat of childhood obesity and undernutrition in low and middle income countries [Report]. World Health Organization. Geneva, Switzerland.
- WHO. (2015). Protein and Amino Acid Requirements in Human Nutrition. World Health Organization. Retrieved from [http://www.who.int/nutrition/publications/nutrientrequirements/WHO\\_TRS\\_935/en/](http://www.who.int/nutrition/publications/nutrientrequirements/WHO_TRS_935/en/).
- WHO. (2020). Increasing fruit and vegetable consumption to reduce risk of noncommunicable diseases [Report]. World Health Organization.
- WHO. (2021). Obesity and overweight. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. [Accessed 2022].
- WHO. (2022). Obesity and overweight. *Fact sheets*: World Health Organization. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.

- WHO/FAO. (2010). Fats and fatty acids in human nutrition [Report]. Food and Agriculture Organization, World Health Organization. Rome, Italy.
- Wilson, S. E. (2012). Marriage, gender and obesity in later life. *Economics & Human Biology*. **10** (4), 431-453.
- Wrzosek, M. and Wisniewska, K. (2018). Early Onset of Obesity and Adult Onset of Obesity as Factors Affecting Patient Characteristics Prior to Bariatric Surgery. *Springer Obesity Surgery*. **28** (12), 3902-3909.
- Wu, G. (2016). Dietary protein intake and human health. *Food & function*. **7** (3), 1251-1265.
- Wu, S., Wang, R., Jiang, A., Ding, Y., Wu, M., Ma, X., Zhao, Y. and He, J. (2014). Abdominal obesity and its association with health-related quality of life in adults: a population-based study in five Chinese cities. *Health and Quality of Life Outcomes*. **12** (1), 1-11.
- Xiaoguang, C., Li, X., Jigyang, W. and Qiyun, C. (2011). Comparisons of body composition prediction accurancy: A study of 2 bioelectric impedance consumer devices in healthy Chinese persons using DXA and MRI as criteria methods. *Journal of Clinical Densitometry*. **14** (4), 458-464.
- Xu, W., Zhang, H., Paillard-Borg, S., Zhu, H., Qi, X. and Rizzuto, D. (2016). Prevalence of overweight and obesity among Chinese adults: role of adiposity indicators and age. *Obesity Facts*. **9** (1), 17-28.
- Yeung, L. D. and Laquatra, I. (2003). " Handbook of Nutrition" (9th ed.). H.J. HEINZ COMPANY. [95-076641].
- Zhang, F., Ye, J., Zhu, X., Wang, L., Gao, P., Shu, G., Jiang, Q. and Wang, S. (2019). Anti-obesity effects of dietary calcium: The evidence and possible mechanisms. *International journal of molecular sciences*. **20** (12), 3072.

## Appendices

### Appendix A

#### INFORMED CONSENT LETTER

Namaste!

I am Samir khulal, a graduate student of BSc. Nutrition and Dietetics in Central Campus of Technology, Dharan and I am here for my dissertation work in Itahari Sub-metropolitan city for the partial fulfillment of my bachelor's degree in Nutrition and Dietetics. The topic for the study is **“RISK FACTOR ASSOCIATED WITH OVERWEIGHT AND OBESITY AMONG REPRODUCTIVE AGED FEMALES RESIDING IN ITAHARI SUB-METROPOLITAN CITY”**.

This study will provide information about the overweight and obesity status and the risk factors associated with it among the study population. During the course of study, it will be required to measure height and weight of the participants followed by the assessment of socio demographic and economic factors, behavioral factors, physical activity, health related factors and dietary factors.

You will be asked some questions and some physical measurements will also be taken with your approve to participate in this study. This study also aims to make you aware about your nutritional status. Your participation in this study will be voluntary. You might have the privilege of not answering the questions if you feel them sensitive. The privacy of your information will be maintained and they will not be misused. I am hopeful that you will be helping me in this study with your participation.

Do you want to get participated in this study?

Yes, I want to be participated in this study and give my consent to take all the required measurements and the answers for this study.

*Signature of participants:*

*Signature of surveyor:*

Date:

Place:

## Appendix B

### Survey Questionnaires

Participant's Code:

Date of Interview (B.S.): \_\_\_\_/\_\_\_\_/\_\_\_\_

#### A. GENERAL INFORMATION

1. Name of participant:

2. Date of Birth (B.S.): \_\_\_\_/\_\_\_\_/\_\_\_\_ Age: ..... Years

3. Address: \_\_\_\_\_ Sub-metropolitan city \_\_\_\_\_ Ward

4. Religion:

i) Hindu    ii) Buddhist    iii) Muslim    iv) Christian    v) Others \_\_\_\_\_

5. Caste Ethnicity

i) Brahmin    ii) Chhetri    iii) Janajati    iv) Dalit    v) Others \_\_\_\_\_

6. Is your father overweight/ obese?

i) Yes

ii. No

8. Is your mother overweight/obese?

i) Yes

ii) No

9. What was your status in early childhood?

i) Normal

ii) Thin

iii) Overweight

iv) Obese

#### B. FAMILY INFORMATION

1. Number of Family members: \_\_\_\_\_

2. Number of female members: \_\_\_\_\_

3. Number of male numbers: \_\_\_\_\_

4. Number of children in family: \_\_\_\_\_

5.Type of Family:    a) Single                      b) Joint

#### C. Socioeconomic status (kuppuswamy'sscale):

1. Marital Status

i) Married    ii) Unmarried    iii) Divorce/ Separated    iv) Widowed

2. Education level

i) Illiterate

- ii) Primary school
  - iii) Secondary school
  - iv) Higher secondary
  - v) Bachelor & above
3. Occupation:
- i) Unemployed
  - ii) Service (Government / private)
  - iii) Daily wage worker
  - iv) Business
  - v) Farmer
  - vii) other
4. Monthly Family Income (Rs):
- a) <30,000
  - b) ≥30,000
5. What was your age when you were married? \_\_\_\_\_
6. Parity:
7. No of still births \_\_\_\_\_

**D ANTHROPOMETRIC MEASUREMENTS**

Indicators	Readings			Mean
Weight (kg)				
Height (kg)				
Waist circumference				
Hip circumference				

**E PHYSICAL ACTIVITY QUESTIONNAIRE (IPAQ-SHORT)**

1. During the last 7 days, on how many days did you do vigorous physical activities (heavy lifting, digging, aerobics, or fast bicycling for more than 10 minutes at a time)?
- i. \_\_\_\_\_ Days per week
  - ii. Don't Know/Not Sure
  - iii. Refused

2. How much time did you usually spend doing vigorous physical activities on one of those days?

i. \_\_\_\_\_ Hours per day \_\_\_\_\_ Minutes per day

ii. Don't Know/Not Sure

iii. Refused

OR/

How much time in total would you spend over the last 7 days doing vigorous physical activities?

i. \_\_\_\_\_ Hours per week \_\_\_\_\_ Minutes per week

ii. Don't Know/Not Sure

iii. Refused

3. During the last 7 days, on how many days did you do moderate physical activities (carrying light loads, bicycling at a regular pace, or double tennis. NO walking)?

i. \_\_\_\_\_ Days per week

ii. Don't Know/Not Sure

iii. Refused

4. How much time did you usually spend doing moderate physical activities on one of those days?

i. \_\_\_\_\_ Hours per day \_\_\_\_\_ Minutes per day

ii. Don't Know/Not Sure

iii. Refused

OR

What is the total amount of time you spent over the last 7 days doing moderate physical activities?

i. \_\_\_\_\_ Hours per week \_\_\_\_\_ Minutes per week

ii. Don't Know/Not Sure

iii. Refused

5. During the last 7 days, on how many days did you walk for at least 10 minutes at a time?

i. \_\_\_\_\_ Days per week

ii. Don't Know/Not Sure

iii. Refused



6. How much time did you usually spend walking on one of those days?

i. \_\_\_\_\_ Hours \_\_\_\_\_ Minutes per day

ii. Don't Know/Not Sure

iii. Refused

OR

What is the total amount of time you spent walking over the last 7 days?

i. \_\_\_\_\_ Hours per week \_\_\_\_\_ Minutes per week

ii. Don't Know/Not Sure

iii. Refused

7. During the last 7 days, how much time did you usually spend sitting on a week day?

i. \_\_\_\_\_ Hours per weekday \_\_\_\_\_ Minutes per weekday

ii. Don't Know/Not Sure

iii. Refused

OR

What is the total amount of time you spent sitting last Wednesday?

i. \_\_\_\_\_ Hours on Wednesday \_\_\_\_\_ Minutes on Wednesday

ii. Don't Know/Not Sure

iii. Refused

## **F. BEHAVIORAL FACTORS**

1. How often do you eat in front of TV?

i) Daily      ii) Twice a week      iii) 3-4 times a week      iv) Never

2. Do you wake at night, get out of bed and eat?

i) Daily

ii) Twice a week

iii) 3-4 times a week

iv) Never

3. Do you smoke or not?

i) Past smoking

ii) Current smoking

iii) non-smoker

4. Do you drink? (alcoholic drink)?

i) Yes

ii) No

5. How frequently do you consume?
- i) Daily
  - ii) Weekly
  - iii) Twice in a week
  - iv) once in a month
  - v) twice in a month
  - vi) others\_\_\_\_\_
6. How much do you drink at a time?
- i) Half glass (tea cup)
  - ii) One glass
  - iii) two or more
7. What type of drink do you consume?
- i) Rakshi/jandh
  - ii) Beer
  - iii) Wine
  - iv)Vodka/Rum
  - v)others\_\_\_\_\_
8. How often do you skip breakfast?
- i) Daily
  - ii) Once a week
  - iii) twice/Thrice a week
  - iv) Never
9. How many hours do you sleep at night?
- i) 4 hours
  - ii) 4-6 hours
  - iii) 8 hours
  - iv) None of them(mention)
10. How often do you have stress?
- i) Daily
  - ii) 2-3 times in a week
  - iii) Never
11. Do you use food as a stress relieving method?
- i) Yes

ii) No

12. If yes which type of food do you prefer?

- i) Processed packet food
- ii) Fruits and vegetables
- iii) Cereals

13. Do you have menstrual irregularities?

- i) Yes
- ii) No

14. How many times do you eat away from house in a day?

- i) Once
- ii) twice
- iii) 2-3 times
- iv) rarely

### G. DIETARY FACTORS

1. What are you, according to your eating pattern?

- i) vegan
- ii) lacto-veg
- iii) lacto-ovo-veg
- iv) non-veg

2. If non-vegetarian which meat do you usually eat?

- i) white (chicken/duck/fish)
- ii) red meat (buff/goat)

3. How frequently do you eat?

- i) Once a week
- ii) Twice a week
- iii) Thrice a week
- iv) Daily

4. How much oil do you use for cooking monthly while cooking? ..... Liters.

5. How many packets of salt do you use monthly? .....

6. How many meals do you eat in a day? .....

### H. FOOD FREQUENCY TABLE

S.N	Type of food	Regular ( at least once a day)	Frequent (3/4 times a week)	Rare (once in a week or less)	Never
<b>1</b>	<b>Cereals</b>				
	Rice				
	Wheat				
	Maize/Millet/Barley				
<b>2</b>	<b>Pulses/Legumes</b>				
	Whole daal				

	Grams/Beans/Peas				
<b>3</b>	<b>Green Leafy Vegetables</b>				
<b>4</b>	<b>Other vegetables</b>				
<b>5</b>	<b>Roots and tubers</b>				
<b>6</b>	<b>Fruits</b>				
<b>7</b>	<b>Dairy products</b>				
	Milk				
	Curd				
	Ghee/Butter				
	Paneer/Cheese				
<b>8</b>	<b>Meat</b>				
	White meat (chicken/fish)				
	Red meat (mutton/beef/pork)				
	egg				
<b>9</b>	<b>Processed Packaged foods/ fast foods</b>				


### I. 24-HOURS DIETARY RECALL

<b>Timing</b>	<b>Description of food</b>	<b>Serving size</b>	<b>Amounts</b>
<b>Breakfast</b>			
<b>Lunch</b>			
<b>Snacks</b>			
<b>Dinner</b>			

Appendix C

Study site

Itahari Sub-metropolitan City



(Municipality, 2079)

P1:Map of Itahari Sub-metropolitan city

## Appendix D

### Photo gallery

